arXiv:1205.5447v2 [math.AP] 3 Jul 2012

HÖLDER ESTIMATES FOR SOLUTIONS OF THE CAUCHY PROBLEM FOR THE POROUS MEDIUM EQUATION WITH EXTERNAL FORCES

MASASHI MIZUNO

ABSTRACT. We study the interior Hölder regularity problem for weak solutions of the porous medium equation with external forces. Since the porous medium equation is the typical example of degenerate parabolic equations, Hölder regularity is a delicate matter and does not follow by classical methods. Caffrelli-Friedman, and Caffarelli-Vazquez-Wolansky showed Hölder regularity for the model equation without external forces. DiBenedetto and Friedman showed the Hölder continuity of weak solutions with some integrability conditions of the external forces but they did not obtain the quantitative estimates. The quantitative estimates are important for studying the perturbation problem of the porous medium equation. We obtain the scale invariant Hölder estimates for weak solutions of the porous medium equations with the external forces. As a particular case, we recover the well known Hölder estimates for the linear heat equation.

1. INTRODUCTION

We consider the following degenerate parabolic equation:

$$\begin{cases} \partial_t u - \Delta u^m = \operatorname{div} f + g, \quad t > 0\,,\, x \in \mathbb{R}^n, \\ \qquad u(0,x) = u_0(x) \geq 0, \quad x \in \mathbb{R}^n, \end{cases} \tag{1.1}$$

where $m > 1$ is a constant, $u = u(t,x) : (0,\infty) \times \mathbb{R}^n \to \mathbb{R}$ is unknown, $u_0 = u_0(x) : \mathbb{R}^n \to [0,\infty)$, $f = f(t,x) : (0,\infty) \times \mathbb{R}^n \to \mathbb{R}^n$ and $g = g(t,x) : (0,\infty) \times \mathbb{R}^n \to \mathbb{R}$ are given. For $f, g \equiv 0$, the equation (1.1) is called the porous medium equation. The equation (1.1) is a degenerate parabolic equation since the diffusion coefficient mu^{m-1} may vanish. It is well-known that solutions of the degenerate parabolic equation (1.1) are not generally smooth even if the initial datum u_0 is smooth enough. We now introduce the notion of weak solutions.

Definition 1.1. For $u_0 \in L^1(\mathbb{R}^n)$ and for $f, g \in L^1((0,\infty) \times \mathbb{R}^n)$, we call u a weak solution of (1.1) if there exists $T > 0$ such that

1. $u(t,x) \geq 0$ for almost all $(t,x) \in [0,T) \times \mathbb{R}^n$;
2. $u \in L^\infty(0,T\,;\, L^1(\mathbb{R}^n) \cap L^{m+1}(\mathbb{R}^n))$ with $\nabla u^m \in L^2((0,T) \times \mathbb{R}^n)$;
3. u satisfies (1.1), namely for all $\varphi \in C^1(0,T\,;\, C^1_0(\mathbb{R}^n))$ and for almost all $0 < t < T$,

$$\begin{aligned} \int_{\mathbb{R}^n} u(t)\varphi(t)\,dx - \int_0^t \int_{\mathbb{R}^n} u\partial_t\varphi\,d\tau dx + \int_0^t \int_{\mathbb{R}^n} \nabla u^m \cdot \nabla\varphi\,d\tau dx \\ = \int_{\mathbb{R}^n} u_0\varphi(0)\,dx - \int_0^t \int_{\mathbb{R}^n} f \cdot \nabla\varphi\,d\tau dx + \int_0^t \int_{\mathbb{R}^n} g\varphi\,d\tau dx. \end{aligned}$$

We remark that the existence of weak solutions is shown by Oleĭnik-Kalašinkov-Čžou [19] and J. L. Lions [16] (cf. Ôtani [22]). Our aim in this paper is to obtain a priori Hölder estimates for weak solutions of (1.1).

Caffarelli-Friedman [5] and Caffarelli-Vázquez-Wolanski [6] showed Hölder continuity for solutions of the porous medium equation. They essentially use a pointwise estimates for the derivative of solutions given by Aronson-Benilan [2] and the comparison principle for the porous medium equation. For the general case with the external force (1.1), the Aronson-Benilan type estimate is not known. In addition, if the equation involves non-local effect such as the system with other equations, the comparison principle does not generally hold. For instance, we consider the following degenerate Keller-Segel system:

$$\begin{cases} \partial_t u - \Delta u^m + \operatorname{div}(u\nabla\psi) = 0, & t > 0,\ x \in \mathbb{R}^n, \\ \qquad -\Delta\psi + \psi = u, & t > 0,\ x \in \mathbb{R}^n, \\ \qquad u(0,x) = u_0(x), & x \in \mathbb{R}^n. \end{cases} \tag{1.2}$$

2000 *Mathematics Subject Classification.* 35K65,35B45,35B65.

It is known by Sugiyama-Kunii [24] that there exists a time global bounded weak solution (u, ψ) of (1.2) in the case of $1 \le m \le 2 - \frac{2}{n}$ and $n \ge 3$ for small initial data. Regularity estimates of solutions of (1.2) are closely related to the large time asymptotic behavior of solutions of (1.2) (cf. Luckhaus-Sugiyama [17], Ogawa [20], Ogawa-Mizuno [21]), however comparison principles do not hold for (1.2). Therefore it is an worth to derive the regularity of the weak solution of (1.1) without using the comparison principle.

On the other hand, DiBenedetto-Friedman [9], Wiegner [26] considered the p-Laplace evolution equation:

$$\begin{cases} \partial_t v - \operatorname{div}(|\nabla v|^{p-2}\nabla v) = 0, & t > 0\,,\, x \in \mathbb{R}^n, \\ \qquad\qquad v(0,x) = v_0(x), & x \in \mathbb{R}^n. \end{cases} \tag{1.3}$$

The p-Laplace evolution equation is a typical example of degenerate parabolic equations. They showed the Hölder continuity for the gradient of the solutions of (1.3) by using the method of alternative and intrinsic rescaling. Misawa [18] showed the gradient Hölder estimates for more general p-Laplace evolution equations. We remark that they does not rely on the comparison principle for the p-Laplace evolution equation (1.3). Roughly speaking, the gradient of the solution may be regarded to satisfy (1.1) with $f, g \equiv 0$ and it seems possible to apply their methods for solutions of (1.1). In fact, DiBenedetto-Friedman [9] showed Hölder continuity for solutions of (1.1) with $f, g \equiv 0$ and $m > 1$. They also mentioned the Hölder continuity of the weak solution of (1.1) involving the external forces $f \in L^q(0,\infty\,;\, L^p(\mathbb{R}^n))$, $g \in L^{\frac{q}{2}}(0,\infty\,;\, L^{\frac{p}{2}}(\mathbb{R}^n))$ with $\frac{2}{q} + \frac{n}{p} < 1$. In this paper we extend the above mentioned results to the case of general external forces, more specifically, we prove Hölder continuity for bounded weak solutions of (1.1). In addition, we obtain Hölder estimates with explicit dependence on the external forces f and g.

It is well-known that Harnack estimates are closely related to Hölder continuity of solutions (cf. Aronson-Caffarelli [3], DiBenedetto [7], [8] and DiBenedetto-Gianazza-Vespri [10], [11]). We remark that the porous medium equation (1.1) is not additive, in particular for a solution u of (1.1) and a constant $k \in \mathbb{R}$, both $u - k$ and $k - u$ do not satisfy (1.1). Thus the Harnack inequality does not imply Hölder continuity of solutions of (1.1) directly. DiBenedetto-Gianazza-Vespri [10], [11] pointed out this fact and considered Hölder estimates for the singular porous medium equation, namely (1.1) with $0 < m < 1$. It is also well-known that regularity of the gradient of solutions imply Hölder continuity. Gradient estimates for p-Laplace evolution equations are recently studied by Kinnunen-Lewis [13], Acerbi-Mingione [1], Duzaar-Mingione [12] and Kuusi-Mingione [14].

Before stating our main theorem, we introduce weak L^p spaces.

Definition 1.2. For a domain $\Omega \subset \mathbb{R}^n$ and an exponent $p > 1$, a function $f \in L^1_{\mathrm{loc}}(\Omega)$ belongs to $L^p_{\mathrm{w}}(\Omega)$ if

$$\|f\|_{L^p_{\mathrm{w}}(\Omega)} := \sup_{K \subset \Omega\,:\,\mathrm{compact}} \frac{1}{|K|^{1-\frac{1}{p}}} \int_K |f|\,dx < \infty.$$

Remark 1.3. By Hölder inequality, we find $L^p(\Omega) \subset L^p_{\mathrm{w}}(\Omega)$. In fact, $L^p_{\mathrm{w}}(\Omega)$ is strictly larger than $L^p(\Omega)$ since $|x|^{-\frac{n}{p}} \notin L^p(\mathbb{R}^n)$ but belonging to $L^p_{\mathrm{w}}(\mathbb{R}^n)$.

Now, we state our main theorem.

Theorem 1.4. *Let $m > 1$ and let u be a bounded weak solution of* (1.1)*. Assume $f \in L^q(0,\infty\,;\, L^p_{\mathrm{w}}(\mathbb{R}^n))$ and $g \in L^{\frac{q}{2}}(0,\infty\,;\, L^{\frac{p}{2}}_{\mathrm{w}}(\mathbb{R}^n))$ for some $p, q > 2$ satisfying $\frac{2}{q} + \frac{n}{p} < 1$. Then, for all $\varepsilon > 0$, the solution u is uniform Hölder continuous with respect to (t,x) in $(\varepsilon,\infty) \times \mathbb{R}^n$. Precisely, there exist constants $C, \sigma > 0$ such that*

$$\begin{aligned} |u(t,x) - u(s,y)| \le C(&\|u\|_{L^\infty((0,\infty)\times\mathbb{R}^n)} \\ &+ \|u\|^{\frac{1}{q}(1-\frac{1}{m})}_{L^\infty((0,\infty)\times\mathbb{R}^n)} \|f\|^{\frac{1}{m}}_{L^q(0,\infty\,;\,L^p_{\mathrm{w}}(\mathbb{R}^n))} + \|u\|^{\frac{2}{q}(1-\frac{1}{m})}_{L^\infty((0,\infty)\times\mathbb{R}^n)} \|g\|^{\frac{1}{m}}_{L^{\frac{q}{2}}(0,\infty\,;\,L^{\frac{p}{2}}_{\mathrm{w}}(\mathbb{R}^n))}) \\ &\times (\|u\|^{\frac{\sigma}{2}(1-\frac{1}{m})}_{L^\infty((0,\infty)\times\mathbb{R}^n)} |t-s|^{\frac{\sigma}{2}} + |x-y|^{\sigma}) \end{aligned} \tag{1.4}$$

for all $(t,x), (s,y) \in (\varepsilon,\infty) \times \mathbb{R}^n$, where $\sigma > 0$ depends only on n, m, p, q and $C > 0$ depends only on n, m, p, q, ε.

When the initial datum u_0 is bounded positive and the external force $\operatorname{div} f + g$ is bounded, then solutions of (1.1) is bounded on some time interval $(0,T)$ by the maximum principle. Sugiyama-Kunii [24] and Ogawa [20] showed the boundedness for the solution and rescaled solution of (1.2) hence we may apply our results for (1.2).

We emphasize that our estimate (1.4) is *scale invariant*. In fact, for the parameter $M > 0$, we consider the scale transform

$$t = \frac{1}{M^{m-1}} s, \quad u_M(s,x) = \frac{1}{M} u(t,x), \tag{1.5}$$
$$f_M(s,x) = f(t,x), \qquad g_M(s,x) = g(t,x).$$

Then u_M satisfies

$$\partial_s u_M - \Delta u_M^m = \operatorname{div}\left(\frac{1}{M^m} f_M\right) + \left(\frac{1}{M^m} g_M\right)$$

and Theorem 1.4 implies

$$\begin{aligned}|u_M(s,y) - u_M(s',y')| \le C\Bigg(&\|u_M\|_{L^\infty((0,\infty)\times\mathbb{R}^n)} \\ &+ \left(\frac{1}{M^m}\right)^{\frac{1}{m}} \|u_M\|_{L^\infty((0,\infty)\times\mathbb{R}^n)}^{\frac{1}{q}(1-\frac{1}{m})} \|f_M\|_{L^q(0,\infty\,;\,L^p_{\mathrm{w}}(\mathbb{R}^n))}^{\frac{1}{m}} \\ &+ \left(\frac{1}{M^m}\right)^{\frac{1}{m}} \|u_M\|_{L^\infty((0,\infty)\times\mathbb{R}^n)}^{\frac{2}{q}(1-\frac{1}{m})} \|g_M\|_{L^{\frac{q}{2}}(0,\infty\,;\,L^{\frac{p}{2}}_{\mathrm{w}}(\mathbb{R}^n))}^{\frac{1}{m}}\Bigg) \\ &\times \left(\|u_M\|_{L^\infty((0,\infty)\times\mathbb{R}^n)}^{\frac{\sigma}{2}(1-\frac{1}{m})} |s-s'|^{\frac{\sigma}{2}} + |y-y'|^\sigma\right).\end{aligned}$$

By the scale transform (1.5), we have (1.4) hence we find that the Hölder estimates (1.4) is invariant for the scale transform (1.5).

Our Hölder estimates (1.4) is some generalization for the case of the heat equation. Actually, letting $m \to 1$, we find that the constant $C > 0$ is bounded and the Hölder exponent $\sigma > 0$ is away from 0. Therefore the estimates (1.4) implies the well-known Hölder estimates for the case of the heat equation.

The basic strategy to prove Theorem 1.4 is to use the method of alternative and intrinsic scaling by DiBenedetto-Friedman [9]. Since they use the local oscillation of solutions as the intrinsic scaling, it seems difficult to obtain Hölder estimates of solutions. On the other hand, we use the local maximum of solutions as the intrinsic scaling and we make the more exact Caccioppoli estimate. The Caccioppoli estimate plays an important role to show the method of alternative. Reconstructing the iteration argument, we obtain the Hölder estimates of solutions.

For an application, we may consider the external force as the perturbation of solutions(cf. Ogawa-Mizuno [21]). Applying our theorem, we do not need L^p integrability of the external force, but growth order of L^2 integral. Therefore, it is useful to study L^2 theory of non-linear degenerate parabolic equations. Furthermore, we can exactly estimate the Hölder norm of solutions by the external force and the maximum of solutions.

The paper is organized as follows. In section 2, we first give an alternative lemma and show Theorem 1.4 using the alternative lemma. The alternative lemma gives either the better lower bounds or the better upper bounds of solutions. We show the lower bounds of the solution in section 3 and the upper bounds of solution in section 4. In appendix, we give some fundamental results of calculus which are necessary for the proof of the main theorem.

At the end of this section, we introduce some notations. For $\rho, M, \theta_0 > 0$ and $t_0 \in \mathbb{R}$, we let open intervals

$$I_\rho(t_0) = (t_0 - \rho^2, t_0), \quad I_\rho^{\theta_0}(t_0) = \left(t_0 - \frac{\theta_0}{2}\rho^2, t_0\right)$$

and

$$I_{\rho,M}(t_0) = \left(t_0 - \frac{\rho^2}{M^{1-\frac{1}{m}}}, t_0\right), \quad I_{\rho,M}^{\theta_0}(t_0) = \left(t_0 - \frac{\theta_0}{2}\frac{\rho^2}{M^{1-\frac{1}{m}}}, t_0\right).$$

For $\rho > 0$ and $x_0 \in \mathbb{R}^n$, we denote the n-dimensional open ball with radius ρ and center x_0 by $B_\rho(x_0)$. We define parabolic cylinders $Q_\rho(t_0,x_0)$, $Q_\rho^{\theta_0}(t_0,x_0)$ and modified parabolic cylinders $Q_{\rho,M}(t_0,x_0)$, $Q_{\rho,M}^{\theta_0}(t_0,x_0)$ by

$$Q_\rho(t_0,x_0) = I_\rho(t_0) \times B_\rho(x_0), \quad Q_\rho^{\theta_0}(t_0,x_0) = I_\rho^{\theta_0}(t_0) \times B_\rho(x_0)$$

and

$$Q_{\rho,M}(t_0,x_0) = I_{\rho,M}(t_0) \times B_\rho(x_0), \quad Q_{\rho,M}^{\theta_0}(t_0,x_0) = I_{\rho,M}^{\theta_0}(t_0) \times B_\rho(x_0).$$

We often abbreviate the center of parabolic cylinders (t_0,x_0). We denote the n-dimensional Lebesgue measure by m_n. A function space $L^q(I_{\rho,M}; L^p_{\mathrm{w}}(B_\rho))$ is abbreviated to $L^q(L^p_{\mathrm{w}})(Q_{\rho,M})$ and another function spaces are also same. For a function f on a set A, we denote the oscillation of f in A by $\operatorname{osc}_A f := \sup_A f - \inf_A f$. We denote the positive part of f and the negative part of f by $f_+ := \max\{0, f\}$ and $f_- := \max\{0, -f\}$, respectively. We



FIGURE 1. the usual parabolic cylinder and the modified parabolic cylinder

remark that a superscript plus or minus are different of the positive part or the negative part. For a constant $k \in \mathbb{R}$ and a function f on a set Ω, we let

$$\{f > k\} := \{x \in \Omega : f(x) > k\}$$

and other level sets such as $\{f < k\}$ are defined in a similar manner. We put $\sigma_0 = 1 - \frac{2}{q} - \frac{n}{p}$ and $h(\rho, M, \omega) := \big\|f\big\|^2_{L^q(L^p_{\mathrm{w}})(Q_{\rho,M})} + \omega \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}$. We denote a constant depending on $m, \beta, \ldots$ by $C(m, \beta, \ldots)$. The same letter C will be used to denote different constants. We use subscript numbers if we consider the relation between the constants. For a open interval $(a, b) \subset \mathbb{R}$ and a open ball $B_\rho(x_0) \subset \mathbb{R}^n$, we call $\eta = \eta(t, x)$ a cut-off function in $Q = (a, b) \times B_\rho(x_0)$ if $\eta \in C^\infty(Q)$ satisfies

$$\eta(t,x) \equiv 0 \quad a \le t \le b, \quad x \in \partial B_\rho(x_0) \quad \text{and} \quad \eta(a,x) \equiv 0 \quad x \in B_\rho(x_0).$$

2. ALTERNATIVE LEMMA AND PROOF OF THE MAIN THEOREM

We hereafter replace u^m by u and we consider the following equation:

$$\partial_t u^{\frac{1}{m}} - \Delta u = -\operatorname{div} f + g. \tag{2.1}$$

Let M and ω be an approximated supremum and oscillation of the weak solution u of (2.1), namely

$$\sup_{Q_{\rho,M}(t_0,x_0)} u \le M \le 3 \sup_{Q_{\rho,M}(t_0,x_0)} u, \tag{2.2}$$

and

$$\frac{3}{4}\omega \le \operatorname*{osc}_{Q_{\rho,M}(t_0,x_0)} u \le \omega. \tag{2.3}$$

Lemma 2.1 (alternative lemma)**.** *Let us assume* (2.2) *and* (2.3)*. Then there exist constants* $0 < \theta_0, \eta_0 < 1$ *and* $\delta_0 > 0$ *depending only on* n, m, p, q *such that for all* $\rho > 0$ *satisfying* $\rho^{\sigma_0} \le \delta_0 \omega M^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-\frac{1}{2}}$*, we obtain the following estimates:*

(i) Lower bounds. If

$$m_{n+1}\left(Q_{\rho,M}(t_0,x_0) \cap \left\{u < \inf_{Q_{\rho,M}(t_0,x_0)} u + \frac{\omega}{2}\right\}\right) \le \theta_0 m_{n+1}\Big(Q_{\rho,M}(t_0,x_0)\Big),$$

then

$$u(t,x) \ge \inf_{Q_{\rho,M}(t_0,x_0)} u + \eta_0 \omega \quad \textit{for } (t,x) \in Q_{\frac{\rho}{2},M}(t_0,x_0);$$

(ii) Upper bounds. If

$$m_{n+1}\left(Q_{\rho,M}(t_0,x_0) \cap \left\{u < \inf_{Q_{\rho,M}(t_0,x_0)} u + \frac{\omega}{2}\right\}\right) > \theta_0 m_{n+1}\Big(Q_{\rho,M}(t_0,x_0)\Big),$$

then

$$u(t,x) \le \sup_{Q_{\rho,M}(t_0,x_0)} u - \eta_0 \omega \quad \textit{for } (t,x) \in Q^{\theta_0}_{\frac{\rho}{2},M}(t_0,x_0).$$

We will prove part (i), which is Proposition 3.1 in Section 3 and part (ii), which is Proposition 4.1 in Section 4. According to Lemma 2.1, we obtain

$$\operatorname*{osc}_{Q^{\theta_0}_{\frac{\rho}{2},M}(t_0,x_0)} u \leq \operatorname*{osc}_{Q_{\rho,M}(t_0,x_0)} u - \eta_0\omega \leq (1-\eta_0)\omega \tag{2.4}$$

provided $\rho^{\sigma_0} \leq \delta_0 \omega M^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-\frac{1}{2}}$. We remark that we may take η_0 as small as we want since we obtain by (2.4)

$$\operatorname*{osc}_{Q^{\theta_0}_{\frac{\rho}{2},M}(t_0,x_0)} u \leq (1-\eta_0)\omega \leq (1-\eta)\omega$$

for any $0 < \eta < \eta_0$.

Remark 2.2. We explain an advantage to use the modified parabolic cylinder. For $\rho \ll 1$ and $M > 0$, we consider

$$\partial_t u^{\frac{1}{m}} - \Delta u = -\operatorname{div} f + g \quad \text{in } Q_{\rho,M}.$$

Introducing the scale transform

$$\begin{aligned} &t = \frac{\rho^2}{M^{1-\frac{1}{m}}} s, && x = \rho y, && \\ &u_{\rho,M}(s,y) = \frac{1}{M} u(t,x), && f_{\rho,M}(s,y) = f(t,x), && g_{\rho,M}(s,y) = g(t,x), \end{aligned}$$

we obtain

$$\partial_s u_{\rho,M}^{\frac{1}{m}} - \Delta_y u_{\rho,M} = -\operatorname{div}\Big(\frac{\rho}{M} f_{\rho,M}\Big) + \frac{\rho^2}{M} g_{\rho,M} \quad \text{in } Q_1. \tag{2.5}$$

Since M can be regarded as the supremum of u on $Q_{\rho,M}$ by the assumption (2.2), we may consider (2.5) as the uniformly parabolic equation. Furthermore, in view of

$$\begin{aligned} \Big\| \frac{\rho}{M} f_{\rho,M} \Big\|^2_{L^q(L^p_{\mathrm{w}})(Q_1)} &= \rho^{2(1-\frac{2}{q}-\frac{n}{p})} M^{-2+\frac{2}{q}(1-\frac{1}{m})} \|f\|^2_{L^q(L^p_{\mathrm{w}})(Q_{\rho,M})}, \\ \Big\| \frac{\rho^2}{M} g_{\rho,M} \Big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_1)} &= \rho^{2(1-\frac{2}{q}-\frac{n}{p})} M^{-1+\frac{2}{q}(1-\frac{1}{m})} \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}, \end{aligned}$$

the inequality $1 - \frac{2}{q} - \frac{n}{p} > 0$ is the sufficient condition to ignore the external force.

We now show Theorem 1.4 by temporary admitting Lemma 2.1. We put $Q = (0,\infty) \times \mathbb{R}^n$, $M_0 = \sup_Q u$ and $\omega_0 = M_0$. Let θ_0, δ_0 and η_0 be as in Lemma 2.1. We choose $0 < \rho_0 < \varepsilon$ satisfying

$$\rho_0^{\sigma_0} \leq \delta_0 \omega_0 M_0^{-\frac{1}{q}(1-\frac{1}{m})} \big(\|f\|^2_{L^q(0,\infty\,;\,L^p_{\mathrm{w}}(\mathbb{R}^n))} + \omega_0 \|g\|_{L^{\frac{q}{2}}(0,\infty\,;\,L^{\frac{p}{2}}_{\mathrm{w}}(\mathbb{R}^n))}\big)^{-\frac{1}{2}}.$$

For $(t_0, x_0) \in (0,\infty) \times \mathbb{R}^n$, We denote $Q_0 = Q_{\rho_0,M_0}(t_0,x_0)$, $\mu_0^+ = \sup_{Q_0} u$ and $\mu_0^- = \inf_{Q_0} u$. Then, we find

$$\begin{cases} \operatorname*{osc}_{Q_0} u \leq \omega_0, \\ \sup_{Q_0} u \leq \sup_{Q} u \leq M_0, \\ \rho_0^{\sigma_0} \leq \delta_0 \omega_0 M_0^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho_0, M_0, \omega_0)^{-\frac{1}{2}}. \end{cases}$$

We choose

$$r_0 := \min\Big\{ (1-\eta_0)^{\frac{1}{\sigma_0}},\ \frac{1}{2}\Big(\frac{1}{3}\Big)^{\frac{1}{2}(1-\frac{1}{m})} \Big(\frac{\theta_0}{2}\Big)^{\frac{1}{2}} \Big\}$$

and choose sequences as follows: For $j \in \mathbb{N}$,

$$\begin{aligned} \omega_j &:= (1-\eta_0)\omega_{j-1}, & \rho_j &:= r_0 \rho_{j-1}, \\ M_j &:= \max\{\mu_{j-1}^+,\ \omega_j\}, & Q_j &:= Q_{\rho_j,M_j}(t_0,x_0), \\ \mu_j^+ &:= \sup_{Q_j} u, & \mu_j^- &:= \inf_{Q_j} u. \end{aligned} \tag{2.6}$$

Using Lemma 2.1, we obtain the following oscillation estimates.

Lemma 2.3. *Let* $\{\omega_j, \rho_j, M_j, Q_j\}_{j=0}^{\infty}$ *is defined by the above* (2.6). *Then for* $0 < \delta_0 < 1$ *defined in Lemma 2.1 and for* $j \in \mathbb{N}$, *we obtain*

$$\begin{cases} \operatorname*{osc}_{Q_j} u \le \omega_j, \\ \sup_{Q_j} u \le \sup_{Q_{j-1}} u \le M_j, \\ \rho_j^{\sigma_0} \le \delta_0 \omega_j M_j^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho_j, M_j, \omega_j)^{-\frac{1}{2}}. \end{cases} \tag{2.7}$$

Proof of Lemma 2.3. By the definition of M_j, we obtain $\sup_{Q_j} u \le \sup_{Q_{j-1}} u \le M_j$. Since $r_0 \le (1-\eta_0)^{\frac{1}{\sigma_0}}$ and definition of ω_j, we find

$$\rho_j^{\sigma_0} \le \delta_0 \omega_j M_j^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho_j, M_j, \omega_j)^{-\frac{1}{2}}.$$

We show $\operatorname{osc}_{Q_j} u \le \omega_j$.

To show $\operatorname{osc}_{Q_j} u \le \omega_j$, we make induction. First we consider the case $j = 1$. Either if $\operatorname{osc}_{Q_0} u \le \frac{3}{4}\omega_0$, then we find $Q_1 \subset Q_0$ since $r_0 \le (\frac{3}{4})^{\frac{1}{2}(1-\frac{1}{m})}$ and

$$\frac{M_1}{M_0} \ge \frac{\omega_1}{M_0} = (1-\eta_0) \ge \frac{3}{4}.$$

For this reason, we obtain

$$\operatorname*{osc}_{Q_1} u \le \operatorname*{osc}_{Q_0} u \le \frac{3}{4}\omega_0 \le (1-\eta_0)\omega_0 = \omega_1.$$

Otherwise, if $\frac{3}{4}\omega_0 \le \operatorname{osc}_{Q_0} u \le \omega_0$, we obtain $M_0 = \omega_0 \le \frac{4}{3}\mu_0^+$. Applying Lemma 2.1, we find

$$\operatorname*{osc}_{Q^{\theta_0}_{\frac{\rho_0}{2}, M_0}(t_0, x_0)} u \le (1-\eta_0)\omega_0.$$

Since $r_0 \le \frac{1}{2}\left(\frac{1}{3}\right)^{\frac{1}{2}(1-\frac{1}{m})}\left(\frac{\theta_0}{2}\right)^{\frac{1}{2}}$, we have $Q_1 \subset Q^{\theta_0}_{\frac{\rho_0}{2}, M_0}(t_0, x_0) \subset Q_0$ and hence

$$\operatorname*{osc}_{Q_1} u \le \operatorname*{osc}_{Q^{\theta_0}_{\frac{\rho_0}{2}, M_0}(t_0, x_0)} u \le (1-\eta_0)\omega_0 = \omega_1.$$

In either case, we obtain (2.7) for $j = 1$. Next we assume (2.7) for $j \le k$ and we show for $j = k+1$ using the following inequality:

$$\mu_{k-1}^+ \le \max\left\{ \frac{3}{2(1-\eta_0)}\omega_k \, , \, 3\mu_k^+ \right\}. \tag{2.8}$$

To show (2.8), we consider the case $\mu_{k-1}^- \le \frac{1}{3}\mu_{k-1}^+$ first. Then

$$\mu_{k-1}^+ \le \operatorname*{osc}_{Q_{k-1}} u + \mu_{k-1}^- \le \omega_{k-1} + \frac{1}{3}\mu_{k-1}^+$$

and hence $\mu_{k-1}^+ \le \frac{3}{2}\omega_{k-1} = \frac{3}{2(1-\eta_0)}\omega_k$. For the other case, namely if $\mu_{k-1}^- > \frac{1}{3}\mu_{k-1}^+$, then we have $\mu_{k-1}^+ < 3\mu_{k-1}^- \le 3\mu_k^- \le 3\mu_k^+$ and we obtain (2.8).

We show (2.7) for $j = k+1$. First we consider the case $\operatorname{osc}_{Q_k} u \le \frac{3}{4}\omega_k$ and we show $Q_{k+1} \subset Q_k$. Either if $M_k = \omega_k$, then

$$\frac{M_{k+1}}{M_k} = \frac{M_{k+1}}{\omega_k} \ge \frac{(1-\eta_0)\omega_k}{\omega_k} = (1-\eta_0) \ge \frac{3}{4}.$$

Since $r_0 \le (\frac{3}{4})^{\frac{1}{2}(1-\frac{1}{m})}$, we obtain $Q_{k+1} \subset Q_k$. Otherwise, if $M_k = \mu_{k-1}^+$, we obtain by (2.8)

$$\begin{aligned} \frac{M_{k+1}}{M_k} = \frac{M_{k+1}}{\mu_{k-1}^+} &\ge \frac{M_{k+1}}{\max\left\{ \frac{3}{2(1-\eta_0)}\omega_k \, , \, 3\mu_k^+ \right\}} \\ &\ge \frac{1}{\max\left\{ \frac{3}{2(1-\eta_0)} \frac{\omega_k}{M_{k+1}} \, , \, \frac{3\mu_k^+}{M_{k+1}} \right\}} \\ &\ge \frac{1}{\max\left\{ \frac{3}{2(1-\eta_0)^2} \, , \, 3 \right\}} \ge \frac{1}{3}. \end{aligned}$$

Since $r_0 \le (\frac{1}{3})^{\frac{1}{2}(1-\frac{1}{m})}$, we have $Q_{k+1} \subset Q_k$. In either case, we have $Q_{k+1} \subset Q_k$ and hence

$$\operatorname*{osc}_{Q_{k+1}} u \le \operatorname*{osc}_{Q_k} u \le \frac{3}{4}\omega_k \le \omega_{k+1.}$$

Second we consider the case $\frac{3}{4}\omega_k \le \operatorname{osc}_{Q_k} u \le \omega_k$. Since $\omega_k \le \frac{4}{3}\mu_k^+$, we obtain

$$\mu_{k-1}^+ \le \max\left\{ \frac{3}{2(1-\eta_0)}\omega_k \, , \, 3\mu_k^+ \right\} \le \max\left\{ \frac{2}{(1-\eta_0)}\mu_k^+ \, , \, 3\mu_k^+ \right\} \le 3\mu_k^+$$

and hence

$$M_k \le \max\left\{ \frac{4}{3}\mu_k^+ \, , \, 3\mu_k^+ \right\} \le 3\mu_k^+.$$

Hence we may apply Lemma 2.1 and we obtain

$$\operatorname*{osc}_{Q^{\theta_0}_{\frac{\rho_k}{2}, M_k}(t_0,x_0)} u \le (1-\eta_0)\omega_k = \omega_{k+1}.$$

Since $r_0 \le \frac{1}{2}\left(\frac{1}{3}\right)^{\frac{1}{2}(1-\frac{1}{m})}\left(\frac{\theta_0}{2}\right)^{\frac{1}{2}}$ and $\frac{M_{k+1}}{M_k} \ge \frac{\mu_k^+}{3\mu_k^+} = \frac{1}{3}$, we have $Q_{k+1} \subset Q^{\theta_0}_{\frac{\rho_k}{2}, M_k}(t_0, x_0)$ and hence

$$\operatorname*{osc}_{Q_{k+1}} u \le \operatorname*{osc}_{Q^{\theta_0}_{\frac{\rho_k}{2}, M_k}(t_0,x_0)} u \le \omega_{k+1}.$$

□

Proof of Theorem 1.4. Remarking that $M_j \ge M_{j+1}$ for $j \in \mathbb{N}$, we have by Lemma 2.3

$$\operatorname*{osc}_{Q_{\rho_j, M_0}(t_0,x_0)} u \le \operatorname*{osc}_{Q_j} u \le \omega_j.$$

We choose $0 < \sigma < 1$ satisfying $r_0^\sigma \ge 1 - \eta_0$. Then we obtain

$$\operatorname*{osc}_{Q_{\rho_j, M_0}(t_0,x_0)} u \le (1-\eta_0)^j \omega_0 = \omega_0 \left(\frac{\rho_j}{\rho_0}\right)^\sigma.$$

For $\rho \le \rho_0$, there exists $k \in \mathbb{N}_0$ such that $\rho_k \le \rho \le \rho_{k-1}$ and hence

$$\operatorname*{osc}_{Q_{\rho, M_0}(t_0,x_0)} u \le \omega_0 \left(\frac{\rho_{k-1}}{\rho_0}\right)^\sigma = \omega_0 r_0^{-\sigma} \left(\frac{\rho_k}{\rho_0}\right)^\sigma \le M_0 r_0^{-\sigma} \left(\frac{\rho}{\rho_0}\right)^\sigma.$$

Taking $\rho_0 > 0$ as

$$\rho_0^{\sigma_0} = \delta_0 \omega_0 M_0^{-\frac{1}{q}(1-\frac{1}{m})} (\|f\|^2_{L^q(L^p_{\mathrm{w}})(Q)} + \omega_0 \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q)})^{-\frac{1}{2}},$$

we find

$$\operatorname*{osc}_{Q_{\rho, M_0}(t_0,x_0)} u \le C M_0^{1-\frac{\sigma}{\sigma_0}} M_0^{\frac{\sigma}{q\sigma_0}(1-\frac{1}{m})} (\|f\|^2_{L^q(L^p_{\mathrm{w}})(Q)} + \omega_0 \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q)})^{\frac{\sigma}{2\sigma_0}} \rho^\sigma \tag{2.9}$$

for $\rho \le \rho_0$ where the constant C depends only on n, m, p and q. Furthermore, if $\rho > \rho_0$, then

$$\operatorname*{osc}_{Q_{\rho, M_0}(t_0,x_0)} u \le M_0 \left(\frac{\rho}{\rho_0}\right)^\sigma \le C M_0^{1-\frac{\sigma}{\sigma_0}} M_0^{\frac{\sigma}{q\sigma_0}(1-\frac{1}{m})} (\|f\|^2_{L^q(L^p_{\mathrm{w}})(Q)} + \omega_0 \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q)})^{\frac{\sigma}{2\sigma_0}} \rho^\sigma.$$

Therefore, we find

$$\begin{aligned}
\operatorname*{osc}_{Q_{\rho, M_0}(t_0,x_0)} u &\le C(M_0 + M_0^{\frac{1}{q}(1-\frac{1}{m})} (\|f\|^2_{L^q(L^p_{\mathrm{w}})(Q)} + M_0 \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q)})^{\frac{1}{2}} \rho^\sigma \\
&\le C(M_0 + M_0^{\frac{1}{q}(1-\frac{1}{m})} \|f\|_{L^q(L^p_{\mathrm{w}})(Q)} + M_0^{\frac{1}{2}+\frac{1}{q}(1-\frac{1}{m})} \|g\|^{\frac{1}{2}}_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q)}) \rho^\sigma \\
&\le C(M_0 + M_0^{\frac{1}{q}(1-\frac{1}{m})} \|f\|_{L^q(L^p_{\mathrm{w}})(Q)} + M_0^{\frac{2}{q}(1-\frac{1}{m})} \|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q)}) \rho^\sigma
\end{aligned}$$

and proof of Theorem 1.4 is complete. □

3. PROOF OF LOWER BOUNDS (I) OF LEMMA 2.1

Without loss of generality, we assume $t_0 = 0$ by using the parallel translation. We omit the center of ball x_0. We hereafter write $\mu^+ = \sup_{Q_{\rho,M}} u,\ \mu^- = \inf_{Q_{\rho,M}} u$.

In this section, we prove Lower bounds in Lemma 2.1. More precisely, we show the following proposition:

Proposition 3.1 (First alternative)**.** *Let* $\rho > 0$ *satisfying*

$$\rho^{\sigma_0} \le \omega M^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-\frac{1}{2}}.$$

Assume inequalities (2.2) *and* (2.3)*. Then there exists* $0 < \theta_0 < 1$ *depending only on* n, m, p, q *such that if*

$$m_{n+1}\Big(Q_{\rho,M} \cap \Big\{u < \mu^- + \frac{\omega}{2}\Big\}\Big) \le \theta_0 m_{n+1}(Q_{\rho,M}),$$

then

$$u(t,x) \ge \mu^- + \frac{\omega}{4} \quad \textit{for } (t,x) \in Q_{\frac{\rho}{2},M}.$$

To show the lower bounds, the following Caccioppoli estimate plays an important role.

Lemma 3.2 (the Caccioppoli estimate for sub-level sets)**.** *Let* $\eta = \eta(t,x)$ *be a cut-off function in* $Q_{\rho,M}$*. For* $\mu^- < k < \mu^- + \frac{1}{2}\omega$*, there exists a constant* $C > 0$ *depending only on* m *such that*

$$\begin{aligned}
&\sup_{t \in I_{\rho,M}} \int_{B_\rho} (u(t)-k)_-^2 \eta^2\, dx + (\mu^+)^{1-\frac{1}{m}} \iint_{Q_{\rho,M}} |\nabla (u-k)_-|^2 \eta^2\, dtdx \\
&\quad \le C\Big\{\omega \iint_{Q_{\rho,M}} (u-k)_- \eta \partial_t \eta\, dtdx + (\mu^+)^{1-\frac{1}{m}} \iint_{Q_{\rho,M}} (u-k)_-^2 |\nabla \eta|^2\, dtdx \\
&\qquad + (\mu^+)^{1-\frac{1}{m}} h(\rho, M, \omega) \Big(\int_{I_{\rho,M}} m_n\Big(B_\rho \cap \{u(t) < k\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})} dt\Big)^{\frac{2}{q'}}\Big\},
\end{aligned} \tag{3.1}$$

where $\frac{1}{2} = \frac{1}{q} + \frac{1}{q'}$.

Proof. Testing a function $-(u-k)_-\eta^2$ in (2.1), we obtain

$$\begin{aligned}
&\frac{1}{m} \iint_{Q_{\rho,M}} \partial_t \Big(\int_0^{(u-k)_-} (k-\xi)^{\frac{1}{m}-1} \xi\, d\xi\Big) \eta^2\, dtdx \\
&\qquad + \iint_{Q_{\rho,M}} \nabla (u-k)_- \cdot \nabla\{(u-k)_- \eta^2\}\, dtdx \\
&\qquad\qquad = -\iint_{Q_{\rho,M}} f \cdot \nabla\{(u-k)_- \eta^2\}\, dtdx - \iint_{Q_{\rho,M}} g (u-k)_- \eta^2\, dtdx.
\end{aligned}$$

By the integration by parts and the Young inequality, we obtain

$$\begin{aligned}
&\frac{1}{m} \sup_{t \in I_{\rho,M}} \int_{B_\rho} \Big(\int_0^{(u(t)-k)_-} (k-\xi)^{\frac{1}{m}-1} \xi\, d\xi\Big) \eta^2(t)\, dx + \frac{1}{4} \iint_{Q_{\rho,M}} |\nabla (u-k)_-|^2 \eta^2\, dtdx \\
&\quad \le \frac{1}{m} \iint_{Q_{\rho,M}} \Big(\int_0^{(u-k)_-} (k-\xi)^{\frac{1}{m}-1} \xi\, d\xi\Big) \partial_t \eta^2\, dtdx \\
&\qquad + 3 \iint_{Q_{\rho,M}} (u-k)_-^2 |\nabla \eta|^2\, dtdx \\
&\qquad + 2 \iint_{Q_{\rho,M} \cap \{u<k\}} |f|^2 \eta^2\, dtdx + \iint_{Q_{\rho,M} \cap \{u<k\}} |g| (u-k)_- \eta^2\, dtdx.
\end{aligned} \tag{3.2}$$

We estimate the 1st term of the left-hand side of (3.2). Since (2.3) and $k \le \mu^- + \frac{\omega}{2} \le \mu^+ - \operatorname{osc}_{Q_{\rho,M}} u + \frac{\omega}{2} \le \mu^+$, we have

$$(k-\xi)^{\frac{1}{m}-1} \ge k^{\frac{1}{m}-1} \ge (\mu^+)^{\frac{1}{m}-1} \quad \text{for } \xi \ge 0$$

and hence

$$
\begin{aligned}
&\frac{1}{2m}\sup_{t\in I_{\rho,M}}\int_{B_\rho}(u(t)-k)_-^2\eta^2(t)\,dx+\frac{1}{4}(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}}|\nabla(u-k)_-|^2\eta^2\,dtdx\\
&\le\frac{1}{m}(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}}\Big(\int_0^{(u-k)_-}(k-\xi)^{\frac{1}{m}-1}\xi\,d\xi\Big)\partial_t\eta^2\,dtdx\\
&\quad+3(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}}(u-k)_-^2|\nabla\eta|^2\,dtdx+2(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}\cap\{u<k\}}|f|^2\eta^2\,dtdx\\
&\quad+(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}\cap\{u<k\}}|g|(u-k)_-\eta^2\,dtdx\\
&=:I_1+I_2+I_3+I_4.
\end{aligned}
\tag{3.3}
$$

We estimate I_3 and I_4. By the definition of the weak L^p space and by the Hölder inequality, we have

$$
\begin{aligned}
\iint_{Q_{\rho,M}\cap\{u<k\}}|f|^2\eta^2\,dtdx&=\int_{I_{\rho,M}}dt\int_{B_\rho\cap\{u(t)<k\}}|f|^2\,dx\\
&\le\int_{I_{\rho,M}}\big\||f(t)|^2\big\|_{L^{\frac{p}{2}}_{\mathrm{w}}(B_\rho)}m_n\Big(B_\rho\cap\{u(t)<k\}\Big)^{1-\frac{2}{p}}\,dt\\
&\le\big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big(\int_{I_{\rho,M}}m_n\Big(B_\rho\cap\{u(t)<k\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})}\,dt\Big)^{\frac{2}{q'}},
\end{aligned}
$$

and

$$
\begin{aligned}
\iint_{Q_{\rho,M}\cap\{u<k\}}|g|(u-k)_-\eta^2\,dtdx&=\frac{\omega}{2}\int_{I_{\rho,M}}dt\int_{B_\rho\cap\{u(t)<k\}}|g|\,dx\\
&\le\frac{\omega}{2}\int_{I_{\rho,M}}\|g(t)\|_{L^{\frac{p}{2}}_{\mathrm{w}}(B_\rho)}m_n\Big(B_\rho\cap\{u(t)<k\}\Big)^{1-\frac{2}{p}}\,dt\\
&\le\frac{\omega}{2}\|g\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big(\int_{I_{\rho,M}}m_n\Big(B_\rho\cap\{u(t)<k\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})}\,dt\Big)^{\frac{2}{q'}}.
\end{aligned}
$$

Therefore

$$
I_3+I_4\le2(\mu^+)^{1-\frac{1}{m}}h(\rho,M,\omega)\Big(\int_{I_{\rho,M}}m_n\Big(B_\rho\cap\{u(t)<k\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})}\,dt\Big)^{\frac{2}{q'}}.
\tag{3.4}
$$

We estimate I_1. Since

$$
\begin{aligned}
\int_0^{(u-k)_-}(k-\xi)^{\frac{1}{m}-1}\xi\,d\xi&\le-m(u-k)_-\int_0^{(u-k)_-}\frac{\partial}{\partial\xi}(k-\xi)^{\frac{1}{m}}\,d\xi\\
&=m(u-k)_-[k^{\frac{1}{m}}-(k-(u-k)_-)^{\frac{1}{m}}],
\end{aligned}
$$

we have

$$
\begin{aligned}
I_1&\le(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}}[k^{\frac{1}{m}}-(k-(u-k)_-)^{\frac{1}{m}}](u-k)_-\partial_t\eta^2\,dtdx\\
&\le(\mu^+)^{1-\frac{1}{m}}\iint_{Q_{\rho,M}}\Big[\Big(\mu^-+\frac{\omega}{2}\Big)^{\frac{1}{m}}-(\mu^-)^{\frac{1}{m}}\Big](u-k)_-\partial_t\eta^2\,dtdx.
\end{aligned}
$$

Either if $\mu^-\le\frac{1}{2}\mu^+$, then $\mu^+\le\omega+\mu^-$ and hence $\mu^+\le2\omega$. Therefore

$$
(\mu^+)^{1-\frac{1}{m}}\Big[\Big(\mu^-+\frac{\omega}{2}\Big)^{\frac{1}{m}}-(\mu^-)^{\frac{1}{m}}\Big]\le(2\omega)^{1-\frac{1}{m}}\Big(\frac{\omega}{2}\Big)^{\frac{1}{m}}\le2^{1-\frac{2}{m}}\omega
$$

and hence

$$
I_1\le C(m)\omega\iint_{Q_{\rho,M}}(u-k)_-\partial_t\eta^2\,dtdx.
$$

Otherwise, if $\mu^- > \frac{1}{2}\mu^+$, then

$$\begin{aligned}\Big(\mu^- + \frac{\omega}{2}\Big)^{\frac{1}{m}} - (\mu^-)^{\frac{1}{m}} &= \int_0^1 \frac{d}{ds}\Big(\mu^- + \frac{\omega}{2}s\Big)^{\frac{1}{m}}\, ds \\ &= \frac{\omega}{2m}\int_0^1 \Big(\mu^- + \frac{\omega}{2}s\Big)^{\frac{1}{m}-1} ds \le \frac{\omega}{2m}(\mu^-)^{\frac{1}{m}-1} \le \frac{\omega}{2m}\Big(\frac{1}{2}\mu^+\Big)^{\frac{1}{m}-1},\end{aligned}$$

and hence

$$(\mu^+)^{1-\frac{1}{m}}\bigg[\Big(\mu^- + \frac{\omega}{2}\Big)^{\frac{1}{m}} - (\mu^-)^{\frac{1}{m}}\bigg] \le \frac{\omega}{2m}(\mu^+)^{1-\frac{1}{m}}\Big(\frac{1}{2}\mu^+\Big)^{\frac{1}{m}-1} \le C(m)\omega.$$

In either case, we obtain

$$I_1 \le C(m)\omega \iint_{Q_{\rho,M}} (u-k)_- \partial_t \eta^2 \, dtdx. \tag{3.5}$$

Substituting (3.4) and (3.5) for (3.3) we obtain (3.1). □

Proof of Proposition 3.1. We consider the scale transform

$$\begin{aligned} s = M^{1-\frac{1}{m}}t, \quad & \tilde{u}(s,x) = u(t,x), \quad \tilde{\eta}(s,x) = \eta(t,x), \\ \tilde{f}(s,x) = f(t,x), \quad & g(s,x) = g(t,x).\end{aligned}$$

and we put $\tilde{h}(\rho,\omega) := \big\|\tilde{f}\big\|^2_{L^q(L^p_{\mathrm{w}})(Q_\rho)} + \omega\|\tilde{g}\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_\rho)}$. We rewrite the Caccioppoli estimate (3.1) as follows:

$$\begin{aligned}&\sup_{s\in I_\rho}\int_{B_\rho}(\tilde{u}(s)-k)^2_-\tilde{\eta}^2(s)\,dx + \frac{(\mu^+)^{1-\frac{1}{m}}}{M^{1-\frac{1}{m}}}\iint_{Q_\rho}|\nabla(\tilde{u}-k)_-|^2\tilde{\eta}^2\,dsdx \\ &\quad\le C(m)\bigg\{\omega\iint_{Q_\rho}(\tilde{u}-k)_-\partial_s\tilde{\eta}^2\,dsdx + \frac{(\mu^+)^{1-\frac{1}{m}}}{M^{1-\frac{1}{m}}}\iint_{Q_\rho}(\tilde{u}-k)^2_-|\nabla\tilde{\eta}|^2\,dsdx \\ &\qquad + \frac{(\mu^+)^{1-\frac{1}{m}}}{M^{1-\frac{1}{m}}}\tilde{h}(\rho,\omega)\bigg(\int_{I_\rho} m_n\big(B_\rho\cap\{\tilde{u}(s)<k\}\big)^{q'(\frac{1}{2}-\frac{1}{p})}\,ds\bigg)^{\frac{2}{q'}}\bigg\}.\end{aligned} \tag{3.6}$$

We take $p_*, q_* > 0$ as

$$\frac{2}{q'} = \frac{2}{q_*}\Big(1+\frac{2\sigma_0}{n}\Big), \quad q'\Big(\frac{1}{2}-\frac{1}{p}\Big) = \frac{q_*}{p_*}.$$

We remark that $\frac{2}{q_*}+\frac{n}{p_*}=\frac{n}{2}$. For $i\in\mathbb{N}$, we take $\rho=\rho_i$, $k=k_i$, $\tilde{\eta}=\tilde{\eta}_i$ satisfying $\tilde{\eta}_i\equiv 1$ on $Q_{\rho_{i+1}}$ and

$$\begin{aligned} k_i &= \mu^- + \frac{1}{4}\omega + \frac{1}{2^{i+1}}\omega, \qquad \rho_i = \frac{1}{2}\rho + \frac{1}{2^{i+1}}\rho, \\ Y_i &:= \frac{m_{n+1}\big(Q_{\rho_i}\cap\{\tilde{u}<k_i\}\big)}{m_{n+1}\big(Q_\rho\big)}, \\ Z_i &= \frac{\rho^2}{m_{n+1}\big(Q_\rho\big)}\bigg(\int_{I_{\rho_i}} m_n\Big(B_{\rho_i}\cap\{\tilde{u}(s)<k_i\}\Big)^{\frac{q_*}{p_*}}ds\bigg)^{\frac{2}{q_*}}, \\ |\nabla\tilde{\eta}_i| &\le \frac{2}{\rho_i-\rho_{i+1}} \le \frac{8\cdot 2^i}{\rho}, \qquad \partial_s\tilde{\eta}_i \le \frac{2}{\rho_i^2-\rho_{i+1}^2} \le \frac{16\cdot 2^{2i}}{3\rho^2}.\end{aligned}$$

Then, by using (2.2) and $(\tilde{u}-k_i)_- \le \frac{\omega}{2}$, we rewrite (3.6) as

$$\begin{aligned}
&\|(\tilde{u}-k_i)_-\tilde{\eta}_i\|^2_{L^\infty(L^2)\cap L^2(H^1)(Q_{\rho_i})} \\
&\le C(m)\Bigg\{\omega\iint_{Q_{\rho_i}}(\tilde{u}-k_i)_-\partial_s\tilde{\eta_i}^2\,dsdx+\iint_{Q_{\rho_i}}(\tilde{u}-k_i)_-^2|\nabla\tilde{\eta}_i|^2\,dsdx \\
&\quad+\tilde{h}(\rho,\omega)\bigg(\int_{I_{\rho_i}}m_n\Big(B_{\rho_i}\cap\{\tilde{u}(s)<k_i\}\Big)^{\frac{q_*}{p_*}}\,ds\bigg)^{\frac{2}{q_*}(1+\frac{2\sigma_0}{n})}\Bigg\} \\
&\le C(m)\Bigg\{\frac{2^{2i}\omega^2}{\rho^2}m_{n+1}\Big(Q_{\rho_i}\cap\{u<k_i\}\Big) \\
&\quad+\tilde{h}(\rho,\omega)\bigg(\int_{I_{\rho_i}}m_{n+1}\Big(B_{\rho_i}\cap\{\tilde{u}(s)<k_i\}\Big)^{\frac{q_*}{p_*}}\,ds\bigg)^{\frac{2}{q_*}(1+\frac{2\sigma_0}{n})}\Bigg\} \\
&\le C(m)\frac{\omega^2 m_{n+1}(Q_\rho)}{\rho^2}\Bigg\{2^{2i}Y_i+\tilde{h}(\rho,\omega)\omega^{-2}\bigg(\frac{m_{n+1}(Q_\rho)}{\rho^2}\bigg)^{\frac{2\sigma_0}{n}}Z_i^{1+\frac{2\sigma_0}{n}}\Bigg\}.
\end{aligned}$$

Using the Ladyženskaja inequality (cf. Proposition A.1) and the Hölder inequality, we have

$$\begin{aligned}
\|(\tilde{u}-k_i)_-\tilde{\eta}_i\|^2_{L^2(Q_{\rho_i})} &\le \|(\tilde{u}-k_i)_-\tilde{\eta}_i\|^2_{L^{2+\frac{4}{n}}(Q_{\rho_i})}\|\chi_{\{\tilde{u}<k_i\}}\|^2_{L^{n+2}(Q_{\rho_i})} \\
&\le C(m,n)\omega^2 m_{n+1}(Q_\rho)Y_i^{\frac{2}{n+2}} \\
&\quad\times\Bigg\{2^{2i}Y_i+\tilde{h}(\rho,\omega)\omega^{-2}\bigg(\frac{m_{n+1}(Q_\rho)}{\rho^2}\bigg)^{\frac{2\sigma_0}{n}}Z_i^{1+\frac{2\sigma_0}{n}}\Bigg\}
\end{aligned}$$

and

$$\|(\tilde{u}-k_i)_-\tilde{\eta}_i\|^2_{L^{q_*}(L^{p_*})(Q_{\rho_i})} \le C(m,n)\frac{\omega^2 m_{n+1}(Q_\rho)}{\rho^2}\Bigg\{2^{2i}Y_i+\tilde{h}(\rho,\omega)\omega^{-2}\bigg(\frac{m_{n+1}(Q_\rho)}{\rho^2}\bigg)^{\frac{2\sigma_0}{n}}Z_i^{1+\frac{2\sigma_0}{n}}\Bigg\}.$$

Since

$$\begin{aligned}
\|(\tilde{u}-k_i)_-\tilde{\eta}_i\|^2_{L^2(Q_{\rho_i})} &\ge \|(\tilde{u}-k_i)_-\|^2_{L^2(Q_{\rho_{i+1}}\cap\{\tilde{u}<k_{i+1}\})} \\
&\ge (k_i-k_{i+1})_-^2 m_{n+1}\Big(Q_{\rho_{i+1}}\cap\{\tilde{u}<k_{i+1}\}\Big) \\
&= \frac{\omega^2}{64\cdot 2^{2i}}m_{n+1}(Q_\rho)Y_{i+1}
\end{aligned}$$

and

$$\begin{aligned}
\|(\tilde{u}-k_i)_-\tilde{\eta}_i\|^2_{L^{q_*}(L^{p_*})(Q_{\rho_i})} &\ge \|(\tilde{u}-k_i)_-\|^2_{L^{q_*}(L^{p_*})(Q_{\rho_{i+1}}\cap\{\tilde{u}<k_{i+1}\})} \\
&\ge (k_i-k_{i+1})_-^2\bigg(\int_{I_{\rho_{i+1}}}m_n\Big(B_{\rho_{i+1}}\cap\{\tilde{u}(s)<k_{i+1}\}\Big)^{\frac{q_*}{p_*}}\,ds\bigg)^{\frac{2}{q_*}} \\
&= \frac{\omega^2}{64\cdot 2^{2i}}\frac{m_{n+1}(Q_\rho)}{\rho^2}Z_{i+1},
\end{aligned}$$

we obtain

$$Y_{i+1} \le C(m,n)\Bigg\{2^{4i}Y_i^{1+\frac{2}{n+2}}+2^{2i}\tilde{h}(\rho,\omega)\omega^{-2}\bigg(\frac{m_{n+1}(Q_\rho)}{\rho^2}\bigg)^{\frac{2\sigma_0}{n}}Y_i^{\frac{2}{n+2}}Z_i^{1+\varepsilon}\Bigg\}$$

and

$$Z_{i+1} \le C(m,n)\Bigg\{2^{4i}Y_i+2^{2i}\tilde{h}(\rho,\omega)\omega^{-2}\bigg(\frac{m_{n+1}(Q_\rho)}{\rho^2}\bigg)^{\frac{2\sigma_0}{n}}Z_i^{1+\varepsilon}\Bigg\}.$$

Either if $q \geq p$, then $\frac{q_*}{p_*} \leq 1$ and we obtain

$$\begin{aligned}Z_0 &= \frac{\rho^2}{m_{n+1}(Q_\rho)}\left(\int_{I_{\rho_0}} m_n\Big(B_{\rho_0} \cap \{\tilde{u}(s) < k_0\}\Big)^{\frac{q_*}{p_*}} ds\right)^{\frac{2}{q_*}} \\ &\leq \frac{\rho^2}{m_{n+1}(Q_\rho)}\left(\int_{I_{\rho_0}} m_n\Big(B_{\rho_0} \cap \{\tilde{u}(s) < k_0\}\Big) ds\right)^{\frac{2}{p_*}} \rho^{\frac{4}{q_*}(1-\frac{q_*}{p_*})} \leq C(n,p,q) Y_0^{\frac{2}{p_*}},\end{aligned} \tag{3.7}$$

by the Hölder inequality. Otherwise, if $q < p$, then

$$\begin{aligned}Z_0 &= \frac{\rho^2}{m_{n+1}(Q_\rho)}\left(\int_{I_{\rho_0}} m_n(B_{\rho_0} \cap \{\tilde{u}(s) < k_0\}) m_n\Big(B_{\rho_0} \cap \{\tilde{u}(s) < k_0\}\Big)^{1-\frac{q_*}{p_*}} ds\right)^{\frac{2}{q_*}} \\ &\leq \frac{\rho^2}{m_{n+1}(Q_\rho)} m_n(B_{\rho_0})^{\frac{2}{p_*}-\frac{2}{q_*}}\left(\int_{I_{\rho_0}} m_n\Big(B_{\rho_0} \cap \{\tilde{u}(s) < k_0\}\Big) ds\right)^{\frac{2}{q_*}} \\ &\leq C(n,p,q) Y_0^{\frac{2}{q_*}}.\end{aligned} \tag{3.8}$$

Therefore, by using $\rho^{\sigma_0} \leq \omega \tilde{h}(\rho, \omega)^{-\frac{1}{2}}$ and Lemma A.4, there exists $0 < \theta_0 = \theta_0(n, m, p, q) < 1$ such that if $Y_0 \leq \theta_0$, then $Y_i \to 0$ as $i \to \infty$, i.e.

$$\tilde{u}(s, x) > \mu^- + \frac{\omega}{4} \quad \text{a.a. } (s, x) \in Q_{\frac{\rho}{2}}.$$

□

4. PROOF OF UPPER BOUNDS (II) OF LEMMA 2.1

In this section, we prove Upper bounds in Lemma 2.1. More precisely we show the following proposition:

Proposition 4.1. *Let* $0 < \theta_0 < 1$. *Assume inequalities* (2.2) *and* (2.3). *Then, there exist* $\eta_1, \delta_1 > 0$ *depending only on* n, m, p, q *and* θ_0 *such that if*

$$\rho^{\sigma_0} \leq \delta_1 \omega M^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-\frac{1}{2}}$$

and

$$m_{n+1}\left(Q_{\rho,M} \cap \Big\{u < \inf_{Q_{\rho,M}} u + \frac{\omega}{2}\Big\}\right) > \theta_0 m_{n+1}(Q_{\rho,M}),$$

then

$$u(t, x) \leq \sup_{Q_{\rho,M}} u - \eta_1 \omega \quad \textit{for } (t, x) \in Q^{\theta_0}_{\frac{\rho}{2},M}.$$

Taking θ_0 as in Proposition 3.1, δ_1, $\eta_1 > 0$ as in Proposition 4.1 and

$$\delta_0 = \min\{1, \delta_1\}, \ \eta_0 = \min\Big\{\frac{1}{4}, \eta_1\Big\},$$

we obtain Lemma 2.1.

To prove Proposition 4.1, we first show measure estimates of sub level sets of some time slice.

Lemma 4.2. *Let* $0 < \theta_0 < 1$. *If*

$$m_{n+1}\left(Q_{\rho,M} \cap \Big\{u < \mu^- + \frac{\omega}{2}\Big\}\right) > \theta_0 m_{n+1}(Q_{\rho,M}), \tag{4.1}$$

then for all $0 < \theta < \theta_0$, *there exists* $-\frac{\rho^2}{M^{1-\frac{1}{m}}} < \tau_0 < -\theta\frac{\rho^2}{M^{1-\frac{1}{m}}}$ *depending only on* θ *and* θ_0 *such that*

$$m_n\left(B_\rho \cap \Big\{u(\tau_0) > \mu^- + \frac{\omega}{2}\Big\}\right) \leq \frac{1-\theta_0}{1-\theta} m_n(B_\rho).$$

Proof. By the change of variable $t = \frac{\rho^2}{M^{1-\frac{1}{m}}}s$, $\tilde{u}(s,x) = u(t,x)$ and (4.1), we obtain

$$\begin{aligned}\int_{-1}^{0} m_n\left(B_\rho \cap \left\{\tilde{u}(s) > \mu^- + \frac{\omega}{2}\right\}\right) ds &= \frac{M^{1-\frac{1}{m}}}{\rho^2} m_{n+1}\left(Q_{\rho,M} \cap \left\{u > \mu^- + \frac{\omega}{2}\right\}\right) \\ &\leq \frac{M^{1-\frac{1}{m}}}{\rho^2}\Bigg(m_{n+1}(Q_{\rho,M}) - m_{n+1}\left(Q_{\rho,M} \cap \left\{u < \mu^- + \frac{\omega}{2}\right\}\right)\Bigg) \\ &< \frac{M^{1-\frac{1}{m}}}{\rho^2}(1-\theta_0)m_{n+1}(Q_{\rho,M}) = (1-\theta_0)m_n(B_\rho).\end{aligned}$$

If $m_n\left(B_\rho \cap \{u(s) > \mu^- + \frac{\omega}{2}\}\right) > \frac{1-\theta_0}{1-\theta} m_n\left(B_\rho\right)$ for all $-1 < s < -\theta$, then

$$\begin{aligned}\int_{-1}^{0} m_n\left(B_\rho \cap \left\{\tilde{u}(s) > \mu^- + \frac{\omega}{2}\right\}\right) ds \geq \int_{-1}^{\theta_0} m_n\left(B_\rho \cap \left\{\tilde{u}(s) > \mu^- + \frac{\omega}{2}\right\}\right) ds \\ \geq (1-\theta_0)m_n(B_\rho),\end{aligned}$$

which is contradiction. □

We next show Bernstein type estimates for the positive part of solutions.

Lemma 4.3. *There exist* $r_0, \delta_2 > 0$ *depending only on* n, m, p, q *and* θ_0 *such that*

$$m_n\left(B_\rho \cap \left\{u(t) > \mu^+ - \frac{\omega}{2^{r_0}}\right\}\right) \leq \left(1 - \left(\frac{\theta_0}{2}\right)^2\right) m_n(B_\rho)$$

for $t \in I^{\theta_0}_{\rho,M}$*, provided* $\rho^{\sigma_0} \leq \delta_2 \omega M^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-\frac{1}{2}}$.

Proof. We rewrite (2.1) as

$$\partial_t u - m u^{1-\frac{1}{m}} \Delta u = -m u^{1-\frac{1}{m}} \operatorname{div} f + m u^{1-\frac{1}{m}} g.$$

Let

$$\psi(\xi) := \log_+\left(\frac{H}{H - (\xi - k)_+ + c}\right),$$

where $k = \mu^- + \frac{\omega}{2}$, $H = \mu^+ - k = \operatorname{osc}_{Q_{\rho,M}} u - \frac{\omega}{2}$, $c = \frac{\omega}{2^{r_0}}$ and $r_0 > 2$ be chosen later. We remark that $\psi, \psi', \psi'' = (\psi')^2 \geq 0$, where $f' = \frac{df}{d\xi}$. We take the cut-off function $\eta = \eta(x)$ as

$$\eta \in C_0^\infty(B_\rho),\ \eta \equiv 1 \text{ on } B_{(1-\sigma)\rho} \quad \text{and } |\nabla \eta| \leq \frac{2}{\sigma\rho},$$

where $\sigma > 0$ will be chosen later. Putting $w = \psi(u)$ and taking the test function $(\psi^2)'(u)\eta^2$ in $(\tau_0, t) \times B_\rho$, where τ_0 will be chosen later, we have

$$\begin{aligned}\frac{1}{2}\int_{B_\rho} w^2\eta^2\,dx\Big|_{\tau_0}^{t} + m\int_{\tau_0}^{t}\int_{B_\rho} (\nabla u \cdot \nabla(u^{1-\frac{1}{m}}(\psi^2)'\eta^2))\,dtdx \\ = m\int_{\tau_0}^{t}\int_{B_\rho} (f \cdot \nabla((u^{1-\frac{1}{m}}(\psi^2)'\eta^2))\,dtdx + m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}} g(\psi^2)'\eta^2\,dtdx.\end{aligned}$$

Since

$$\nabla((u^{1-\frac{1}{m}}(\psi^2)'\eta^2) = \left(1 - \frac{1}{m}\right)u^{-\frac{1}{m}}(\psi^2)'\eta^2\nabla u + u^{1-\frac{1}{m}}(\psi^2)''\eta^2\nabla u + u^{1-\frac{1}{m}}(\psi^2)'\nabla\eta^2,$$

we obtain

$$\begin{aligned}
&\frac{1}{2}\int_{B_\rho} w^2\eta^2\,dx\Big|_{\tau_0}^{t} + (m-1)\int_{\tau_0}^{t}\int_{B_\rho} u^{-\frac{1}{m}}(\psi^2)'|\nabla u|^2\eta^2\,dtdx + m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)''|\nabla u|^2\eta^2\,dtdx\\
&= -m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)'(\nabla u\cdot\nabla\eta^2)\,dtdx + (m-1)\int_{\tau_0}^{t}\int_{B_\rho} u^{-\frac{1}{m}}(g^2)'(f\cdot\nabla u)\eta^2\,dtdx\\
&\quad + m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)''(f\cdot\nabla u)\eta^2\,dtdx + m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)'(f\cdot\nabla\eta^2)\,dtdx\\
&\quad + m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)'g\eta^2\,dtdx\\
&=: I_1+I_2+I_3+I_4+I_5.
\end{aligned}$$

Using the property $(\psi^2)'\nabla u = 2w\nabla w$ and the Young inequality, we have

$$\begin{aligned}
I_1 &\le m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w|\nabla w|^2\eta^2\,dtdx + 4m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w|\nabla\eta|^2\,dtdx,\\
I_2 &\le \frac{m-1}{2}\int_{\tau_0}^{t}\int_{B_\rho} u^{-\frac{1}{m}}(\psi^2)'|\nabla u|^2\eta^2\,dtdx + \frac{m-1}{2}\int_{\tau_0}^{t}\int_{B_\rho} u^{-\frac{1}{m}}(\psi^2)'|f|^2\eta^2\,dtdx,\\
I_3 &\le \frac{m}{4}\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)''|\nabla u|^2\eta^2\,dtdx + m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)''|f|^2\eta^2\,dtdx,\\
I_4 &\le 4m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w\psi'|f||\nabla\eta|\eta\,dtdx\\
&\le 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w|\nabla\eta|^2\,dtdx + 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(g')^2w|f|^2\eta^2\,dtdx,\\
I_5 &\le 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w\psi'|g|\eta^2\,dtdx.
\end{aligned}$$

Since $\psi'' = (\psi')^2$, $(\psi^2)'' = 2(\psi')^2(1+\psi)$, we have

$$\begin{aligned}
&m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi^2)''|\nabla u|^2\eta^2\,dtdx\\
&\qquad\qquad = 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}|\nabla w|^2\eta^2\,dtdx + 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w|\nabla w|^2\eta^2\,dtdx.
\end{aligned}$$

Combining the above estimates, we have

$$\begin{aligned}
&\frac{1}{2}\int_{B_\rho} w^2(t)\eta^2(t)\,dx + \frac{m-1}{2}\int_{\tau_0}^{t}\int_{B_\rho} u^{-\frac{1}{m}}(\psi^2)'\eta^2|\nabla u|^2\,dtdx\\
&\quad + \frac{3}{2}m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}|\nabla w|^2\eta^2\,dtdx + \frac{m}{2}\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w|\nabla w|^2\eta^2\,dtdx\\
&\le \frac{1}{2}\int_{B_\rho} w^2(\tau_0)\eta^2(\tau_0)\,dx + 6m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w|\nabla\eta|^2\,dtdx\\
&\quad + \frac{m-1}{2}\int_{\tau_0}^{t}\int_{B_\rho} u^{-\frac{1}{m}}(\psi^2)'|f|^2\eta^2\,dtdx + 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}(\psi')^2(1+2w)|f|^2\eta^2\,dtdx\\
&\quad + 2m\int_{\tau_0}^{t}\int_{B_\rho} u^{1-\frac{1}{m}}w\psi'|g|\eta^2\,dtdx\\
&=: I_6+I_7+I_8+I_9+I_{10}.
\end{aligned} \tag{4.2}$$

For simplicity, we put $k' = \mu^+ - c = \mu^+ - \frac{\omega}{2^{r_0}}$. First, we estimate the left-hand side of (4.2). Since $k' > k$, we have

$$
\begin{aligned}
\frac{1}{2}\int_{B_\rho} w^2(t)\eta^2(t)\,dx &\geq \frac{1}{2}\int_{B_{(1-\sigma)\rho}\cap\{u(t)>k'\}} w^2(t)\,dx \\
&\geq \frac{1}{2}\int_{B_{(1-\sigma)\rho}\cap\{u(t)>k'\}} \log^2\Big(\frac{H}{H-(k'-k)+c}\Big)\,dx \\
&\geq \frac{1}{2}\log^2\Big(\frac{\frac{\omega}{4}}{\frac{\omega}{2^{r_0-1}}}\Big)m_n\Big(B_{(1-\sigma)\rho}\cap\{u(t)>k'\}\Big) \\
&= \frac{1}{2}(r_0-3)^2\log^2 2 m_n\Big(B_{(1-\sigma)\rho}\cap\{u(t)>k'\}\Big).
\end{aligned} \tag{4.3}
$$

Second, we estimate I_6. Taking τ_0 as in Lemma 4.2 with $\theta = \frac{\theta_0}{2}$, we obtain

$$
w = \log_+\Big(\frac{H}{H-(u-k)_+ + c}\Big) \leq \log\Big(\frac{\frac{1}{2}\omega}{\frac{1}{2^{r_0}}\omega}\Big) = (r_0-1)\log 2
$$

and hence

$$
\begin{aligned}
I_6 &\leq \frac{1}{2}\int_{B_\rho\cap\{u(\tau_0)>k\}} w^2(\tau_0)\,dx \\
&\leq \frac{1}{2}(r_0-1)^2\log^2 2|B_\rho\cap\{u(\tau_0)>k\}| \leq \frac{1}{2}\cdot\frac{1-\theta_0}{1-\frac{\theta_0}{2}}(r_0-1)^2\log^2 2 m_n\big(B_\rho\big).
\end{aligned} \tag{4.4}
$$

We estimate I_7. From $t-\tau_0 \leq \frac{\rho^2}{M^{1-\frac{1}{m}}}$ and (2.2), we have

$$
\begin{aligned}
I_7 &\leq 6m(\mu^+)^{1-\frac{1}{m}}(t-\tau_0)(r_0-1)\log 2\Big(\frac{2}{\sigma\rho}\Big)^2 m_n\big(B_\rho\big) \\
&\leq C(m)\Big(\frac{r_0-1}{\sigma^2}\Big)m_n\big(B_\rho\big).
\end{aligned} \tag{4.5}
$$

We estimate I_8. Since

$$
\psi' \leq \frac{1}{H-(u-k)_+ + c} \leq \frac{1}{c} = \frac{2^{r_0}}{\omega}, \quad (\psi^2)' = 2\psi\psi' \leq \frac{2^{r_0+1}}{\omega}(r_0-1)\log 2
$$

and

$$
u^{-\frac{1}{m}} \leq k^{-\frac{1}{m}} \leq \Big(\frac{\omega}{2}\Big)^{-\frac{1}{m}} \quad \text{for } u \geq k,
$$

we have

$$
I_8 \leq C(m)(r_0-1)2^{r_0}\omega^{-1-\frac{1}{m}}\int_{\tau_0}^{t}\int_{B_\rho\cap\{u(s)>k\}} |f|^2\,dtdx.
$$

By the definition of the weak L^p space and by the Hölder inequality, we have

$$
\begin{aligned}
\int_{\tau_0}^{t}\int_{B_\rho\cap\{u(s)>k\}} |f|^2\,dtdx &\leq \int_{\tau_0}^{t}\big\||f(s)|^2\big\|_{L^{\frac{p}{2}}_{\mathrm{w}}(B_\rho)} m_n\Big(B_\rho\cap\{u(s)>k\}\Big)^{1-\frac{2}{p}}\,ds \\
&\leq C(n,p)M^{\frac{2}{q}(1-\frac{1}{m})}\big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\rho^{2\sigma_0}\frac{m_n\big(B_\rho\big)}{M^{1-\frac{1}{m}}}.
\end{aligned}
$$

Using (2.3), we obtain

$$
\begin{aligned}
I_8 &\leq C(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega^2}M^{\frac{2}{q}(1-\frac{1}{m})}\big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big) \\
&\quad\times\Big(\frac{\omega}{M}\Big)^{1-\frac{1}{m}}(r_0-1)2^{r_0}m_n\big(B_\rho\big) \\
&\leq C(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega^2}M^{\frac{2}{q}(1-\frac{1}{m})}\big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big)(r_0-1)2^{r_0}m_n\big(B_\rho\big).
\end{aligned} \tag{4.6}
$$

We estimate I_9 and I_{10}. Considering the same calculation for I_8, we have

$$I_9 \le C(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega^2} M^{\frac{2}{q}(1-\frac{1}{m})} \big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big) 2^{2r_0}(1+2(r_0-1)\log 2) m_n\big(B_\rho\big) \tag{4.7}$$

and

$$I_{10} \le C(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega} M^{\frac{2}{q}(1-\frac{1}{m})} \big\|g\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big) 2^{r_0}(r_0-1) m_n\big(B_\rho\big). \tag{4.8}$$

Combining estimates (4.3)–(4.8), we have

$$\begin{aligned} m_n\Big(B_{(1-\sigma)\rho} \cap \{u(t) > k'\}\Big) \le \Bigg\{ &\frac{1-\theta_0}{1-\frac{\theta_0}{2}}\Big(\frac{r_0-1}{r_0-3}\Big)^2 + \frac{C_1(m)}{\sigma^2}\frac{r_0-1}{(r_0-3)^2} \\ &+ C_2(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega^2} M^{\frac{2}{q}(1-\frac{1}{m})} \big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big) \frac{2^{r_0}(r_0-1)}{(r_0-3)^2} \\ &+ C_3(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega^2} M^{\frac{2}{q}(1-\frac{1}{m})} \big\||f|^2\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big) \frac{2^{2r_0}(1+2(r_0-1)\log 2)}{(r_0-3)^2} \\ &+ C_4(n,m,p)\Big(\frac{\rho^{2\sigma_0}}{\omega} M^{\frac{2}{q}(1-\frac{1}{m})} \big\|g\big\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_{\rho,M})}\Big) \frac{2^{r_0}(r_0-1)}{(r_0-3)^2} \Bigg\} m_n\big(B_\rho\big). \end{aligned}$$

Since

$$\begin{aligned} m_n\big(B_\rho \cap \{u(t)>k'\}\big) &= m_n\Big((B_\rho \setminus B_{(1-\sigma)\rho}) \cap \{u(t)>k'\}\Big) + m_n\Big(B_{(1-\sigma)\rho} \cap \{u(t)>k'\}\Big) \\ &\le (1-(1-\sigma)^n) m_n\big(B_\rho\big) + m_n\Big(B_{(1-\sigma)\rho} \cap \{u(t)>k'\}\Big), \end{aligned}$$

we have

$$\begin{aligned} &m_n\Big(B_{(1-\sigma)\rho} \cap \{u(t) > k'\}\Big) \\ &\quad \le \Bigg\{ \frac{1-\theta_0}{1-\frac{\theta_0}{2}}\Big(\frac{r_0-1}{r_0-3}\Big)^2 + \frac{C_1(m)}{\sigma^2}\frac{r_0-1}{(r_0-3)^2} + (1-(1-\sigma)^n) \\ &\qquad + \max\{C_2, C_3, C_4\} \frac{\rho^{2\sigma_0}}{\omega^2} M^{\frac{2}{q}(1-\frac{1}{m})} h(\rho, M, \omega) C_5(r_0) \Bigg\} m_n\big(B_\rho\big), \end{aligned}$$

where

$$C_5(r_0) = \max\bigg\{ \frac{2^{r_0}(r_0-1)}{(r_0-3)^2}, \frac{2^{2r_0}(1+2(r_0-1)\log 2)}{(r_0-3)^2} \bigg\}.$$

We choose parameters r_0, σ and δ_2. First we choose $\sigma = \sigma(n, \theta_0)$ satisfying $1-(1-\sigma)^n \le \frac{1}{8}\theta_0^2$. Second, we choose $r_0 = r_0(n, m, \theta_0)$ satisfying

$$\Big(\frac{r_0-1}{r_0-3}\Big)^2 \le \Big(1 - \frac{\theta_0}{2}\Big)(1+\theta_0) \ \text{ and } \ \frac{C_1(m)}{\sigma^2}\frac{r_0-1}{(r_0-3)^2} \le \frac{1}{8}\theta_0^2.$$

Finally, we choose $\delta_2 = \delta_2(n, m, p, \theta_0) > 0$ sufficiently small such that

$$\max\{C_2, C_3, C_4\} C_5(r_0) \delta_2 \le \frac{1}{2}\theta_0^2.$$

Then, if $\rho^{2\sigma_0} \le \delta_2 \omega^2 M^{-\frac{2}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-1}$, we have

$$m_n\Big(B_\rho \cap \{u(t) > k'\}\Big) \le \bigg(1 - \Big(\frac{\theta_0}{2}\Big)^2\bigg) m_n\big(B_\rho\big).$$

□

In the proof of Proposition 4.1, we need to show the Caccioppoli estimate.

Lemma 4.4 (the Caccioppoli estimate for super level sets)**.** *Let* $\eta = \eta(t,x)$ *be a cut-off function in* $Q^{\theta_0}_{\rho,M}$*. For* $k \geq \mu^+ - \frac{\omega}{2}$*, there exists a constant* $C > 0$ *depending only on* m *such that*

$$
\begin{aligned}
\sup_{t \in I^{\theta_0}_{\rho,M}} \int_{B_\rho} (u(t)-k)_+^2 \eta^2(t)\, dx + M^{1-\frac{1}{m}} \iint_{Q^{\theta_0}_{\rho,M}} |\nabla (u-k)_+|^2 \eta^2 \, dtdx \\
\leq C \Big\{ \Big(\frac{M}{\mu^+} \Big)^{1-\frac{1}{m}} \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 \partial_t \eta^2 \, dtdx + M^{1-\frac{1}{m}} \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 |\nabla \eta|^2 \, dtdx \\
+ M^{1-\frac{1}{m}} h(\rho, M, \omega) \Big(\int_{t \in I^{\theta_0}_{\rho,M}} m_n \Big(B_\rho \cap \{u(t) > k\} \Big)^{q'(\frac{1}{2}-\frac{1}{p})} dt \Big)^{\frac{2}{q'}} \Big\},
\end{aligned}
\tag{4.9}
$$

where $\frac{1}{2} = \frac{1}{q} + \frac{1}{q'}$*.*

Proof. Testing a function $(u-k)_+\eta^2$ to (2.1), we have

$$
\begin{aligned}
&\frac{1}{m} \sup_{t \in I^{\theta_0}_{\rho,M}} \int_{B_\rho} \Big(\int_0^{(u(t)-k)_+} (k+\xi)^{\frac{1}{m}-1} \xi \, d\xi \Big) \eta^2(t)\, dx + \iint_{Q^{\theta_0}_{\rho,M}} |\nabla (u-k)_+|^2 \eta^2 \, dtdx \\
&\leq \frac{1}{m} \iint_{Q^{\theta_0}_{\rho,M}} \Big(\int_0^{(u-k)_+} (k+\xi)^{\frac{1}{m}-1} \xi \, d\xi \Big) \partial_t \eta^2 \, dtdx - \iint_{Q^{\theta_0}_{\rho,M}} (\nabla (u-k)_+ \cdot \nabla \eta^2)(u-k)_+ \, dtdx \\
&\quad + \iint_{Q^{\theta_0}_{\rho,M}} f \cdot \nabla (u-k)_+ \eta^2 \, dtdx + \iint_{Q^{\theta_0}_{\rho,M}} (f \cdot \nabla \eta^2)(u-k)_+ \, dtdx \\
&\quad + \iint_{Q^{\theta_0}_{\rho,M}} g(u-k)_+ \eta^2 \, dtdx \\
&=: I_1 + I_2 + I_3 + I_4 + I_5.
\end{aligned}
\tag{4.10}
$$

By the Young inequality and since $k > \mu^+ - \frac{\omega}{2}$, we have

$$
\begin{aligned}
I_2 &\leq \frac{1}{2} \iint_{Q^{\theta_0}_{\rho,M}} |\nabla (u-k)_+|^2 \eta^2 \, dtdx + 2 \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 |\nabla \eta|^2 \, dtdx, \\
I_3 &\leq \frac{1}{4} \iint_{Q^{\theta_0}_{\rho,M}} |\nabla (u-k)_+|^2 \eta^2 \, dtdx + \iint_{Q^{\theta_0}_{\rho,M} \cap \{u>k\}} |f|^2 \eta^2 \, dtdx, \\
I_4 &\leq \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 |\nabla \eta|^2 \, dtdx + \iint_{Q^{\theta_0}_{\rho,M} \cap \{u>k\}} |f|^2 \eta^2 \, dtdx, \\
I_5 &\leq \frac{\omega}{2} \iint_{Q^{\theta_0}_{\rho,M} \cap \{u>k\}} |g| \eta^2 \, dtdx.
\end{aligned}
\tag{4.11}
$$

We estimate the first term of the left-hand side in (4.10). Since

$$
(k+\xi)^{\frac{1}{m}-1} \geq u^{\frac{1}{m}-1} \geq (\mu^+)^{\frac{1}{m}-1} \geq M^{\frac{1}{m}-1} \quad \text{for} \quad 0 \leq \xi \leq (u-k)_+,
$$

we have

$$
\int_0^{(u(t)-k)_+} (k+\xi)^{\frac{1}{m}-1} \xi \, d\xi \geq \frac{1}{2} M^{\frac{1}{m}-1} (u(t)-k)_+^2.
\tag{4.12}
$$

Finally, we estimate I_1. By (2.3), we have

$$
(k+\xi)^{\frac{1}{m}-1} \leq k^{\frac{1}{m}-1} \leq \Big(\mu^+ - \frac{\omega}{2} \Big)^{\frac{1}{m}-1} \leq \Big(\frac{1}{3} \mu^+ \Big)^{\frac{1}{m}-1}
$$

and hence

$$
I_1 \leq \frac{1}{2} \Big(\frac{1}{3} \mu^+ \Big)^{\frac{1}{m}-1} \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 \partial_t \eta^2 \, dtdx.
\tag{4.13}
$$

Combining estimates (4.11), (4.12) and (4.13), we obtain

$$
\begin{aligned}
M^{\frac{1}{m}-1} \sup_{t\in I^{\theta_0}_{\rho,M}} \int_{B_\rho} (u(t)-k)_+^2 \eta^2(t)\,dx + \iint_{Q^{\theta_0}_{\rho,\omega}} |\nabla (u-k)_+|^2 \eta^2 \,dtdx \\
\leq C(m)\Big\{ (\mu^+)^{\frac{1}{m}-1} \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 \partial_t \eta^2 \,dtdx + \iint_{Q^{\theta_0}_{\rho,M}} (u-k)_+^2 |\nabla \eta|^2 \,dtdx \\
+ \iint_{Q^{\theta_0}_{\rho,M}\cap\{u>k\}} |f|^2\eta^2\,dtdx + \frac{\omega}{2}\iint_{Q^{\theta_0}_{\rho,M}\cap\{u>k\}} |g|\eta^2\,dtdx \Big\}.
\end{aligned}
\tag{4.14}
$$

Using the same argument of the proof of Lemma 3.2, we have

$$
\iint_{Q^{\theta_0}_{\rho,M}\cap\{u>k\}} |f|^2\eta^2\,dtdx \leq \big\| |f|^2 \big\|_{L^{\frac{q}{2}} L^{\frac{p}{2}}_{\mathrm{w}}(Q^{\theta_0}_{\rho,M})} \left(\int_{I^{\theta_0}_{\rho,M}} m_n\Big(B_\rho \cap \{u(t)>k\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})} \,dt \right)^{\frac{2}{q'}},
$$

and

$$
\iint_{Q^{\theta_0}_{\rho,M}\cap\{u>k\}} |g|\eta^2\,dtdx \leq \| g \|_{L^{\frac{q}{2}} L^{\frac{p}{2}}_{\mathrm{w}}(Q^{\theta_0}_{\rho,M})} \left(\int_{I^{\theta_0}_{\rho,M}} m_n\Big(B_\rho \cap \{u(t)>k\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})} \,dt \right)^{\frac{2}{q'}},
$$

hence we obtain (4.9) from (4.14). □

Using Bernstein type estimates, the Caccioppoli estimate and the hole filling argument, we may prove the smallness of measures of super level sets.

Lemma 4.5. *Let* $\rho_0 = \frac{3}{4}\rho$*. For* $0<\nu<1$*, there exist* q_0 , $\delta_1 > 0$ *depending only on* n, m, p, q, θ_0 *and* ν *such that*

$$
m_{n+1}\left(Q^{\theta_0}_{\rho_0,M} \cap \left\{ u > \mu^+ - \frac{\omega}{2^{q_0+1}} \right\} \right) \leq \nu m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)
$$

provided $\rho^{\sigma_0} \leq \delta_2 \omega M^{-\frac{1}{q}(1-\frac{1}{m})} h(\rho, M, \omega)$.

Remark 4.6. We obtain the estimate of δ_1 as

$$
\delta_1 \leq \theta_0^{\frac{1}{q}-\frac{1}{2}} 2^{-q_0}.
$$

Proof of Lemma 4.5. We fix $t \in I^{\theta_0}_{\rho,M}$ and set

$$
l := \mu^+ - \frac{\omega}{2^{j+1}}, \quad k := \mu^+ - \frac{\omega}{2^j},
$$

where $j \geq r_0$ and the constant r_0 is given by Lemma 4.3. By the Poincaré type inequality (cf. Proposition A.2), we have

$$
\frac{\omega}{2^{j+1}} m_n\big(B_{\rho_0} \cap \{u(t) > l\}\big) \leq \frac{C(n)\rho_0^{n+1}}{m_n\big(B_{\rho_0}\cap\{u(t)\leq k\}\big)} \int_{B_{\rho_0}\cap\{k<u(t)\leq l\}} |\nabla u(t)|\,dx.
$$

Since $k > \mu^+ - \frac{\omega}{2^{r_0}}$ and Lemma 4.3, we have

$$
m_n\big(B_{\rho_0}\cap\{u(t)\leq k\}\big) = m_n\big(B_{\rho_0}\big) - m_n\big(B_{\rho_0}\cap\{u(t)>k\}\big) \geq \left(\frac{\theta_0}{2}\right)^2 m_n\big(B_{\rho_0}\big)
$$

and hence

$$
\frac{\omega}{2^{j+1}} m_n\big(B_{\rho_0}\cap\{u(t)>l\}\big) \leq \frac{C(n)\rho_0}{\theta_0^2} \int_{B_{\rho_0}\cap\{k<u(t)\leq l\}} |\nabla u(t)|\,dx. \tag{4.15}
$$

Integrating over $I^{\theta_0}_{\rho_0,M}$ for (4.15), we obtain

$$
\begin{aligned}
\frac{\omega}{2^{j+1}} m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\cap\{u>l\}\big) &\leq \frac{C(n)\rho_0}{\theta_0^2} \int_{I^{\theta_0}_{\rho_0,M}} \int_{B_{\rho_0}\cap\{k<u(t)\leq l\}} |\nabla u(t)|\,dtdx \\
&\leq \frac{C(n)\rho_0}{\theta_0^2} \|\nabla(u-k)_+\|_{L^2(Q^{\theta_0}_{\rho_0,M})} m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\cap\{k<u\leq l\}\big)^{\frac{1}{2}}.
\end{aligned}
$$

We estimate $\|\nabla(u-k)_+\|_{L^2(Q^{\theta_0}_{\rho_0,M})}$. Let $\eta=\eta(t,x)$ be a cut-off function in $Q^{\theta_0}_{\rho,M}$ satisfying

$$\eta\equiv 1 \text{ on } Q^{\theta_0}_{\rho_0,M},\quad |\nabla\eta|\le\frac{8}{\rho}\quad\text{and}\quad \partial_t\eta\le\frac{10M^{1-\frac{1}{m}}}{\theta_0\rho^2}.$$

Then, by the Caccioppoli estimate (Lemma 4.4), we have

$$\begin{aligned}\|\nabla(u-k)_+\|^2_{L^2(Q^{\theta_0}_{\rho_0,M})} &\le \|\nabla(u-k)_+\eta\|^2_{L^2(Q^{\theta_0}_{\rho,M})}\\ &\le C(m)\Bigg\{\iint_{Q^{\theta_0}_{\rho,M}}(u-k)^2_+\big(|\nabla\eta|^2+(\mu^+)^{\frac{1}{m}-1}\partial_t\eta^2\big)\,dtdx\\ &\qquad+h(\rho,M,\omega)\bigg(\int_{I^{\theta_0}_{\rho,M}}m_n\big(B_\rho\cap\{u(t)>k\}\big)^{q'(\frac{1}{2}-\frac{1}{p})}\,dt\bigg)^{\frac{2}{q'}}\Bigg\}\\ &=: I_1+I_2.\end{aligned}\tag{4.16}$$

First we estimate I_1. By the inequality (2.2), we have

$$\begin{aligned}I_1&\le C(m)(\mu^+-k)^2_+\bigg(\frac{1}{\rho^2}+\frac{M^{1-\frac{1}{m}}}{\theta_0\rho^2}(\mu^+)^{\frac{1}{m}-1}\bigg)m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\\ &\le C(m)\Big(\frac{\omega}{2^j}\Big)^2\frac{1}{\theta_0\rho^2}\Big(\frac{M}{\mu^+}\Big)^{1-\frac{1}{m}}m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\\ &\le C(m)\Big(\frac{\omega}{2^j}\Big)^2\frac{1}{\theta_0\rho^2}m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big).\end{aligned}\tag{4.17}$$

We estimate I_2. Since

$$\begin{aligned}&\bigg(\int_{I^{\theta_0}_{\rho,M}}m_n\big(B_\rho\cap\{u(t)>k\}\big)^{q'(\frac{1}{2}-\frac{1}{p})}\,dt\bigg)^{\frac{2}{q'}}\\ &\le m_n\big(B_\rho\big)^{1-\frac{2}{p}}\bigg(\frac{\theta_0}{2}\frac{\rho^2}{M^{1-\frac{1}{m}}}\bigg)^{\frac{2}{q'}}\\ &\le C(q)m_n\big(B_\rho\big)^{-\frac{2}{p}}\bigg(\frac{\theta_0}{2}\frac{\rho^2}{M^{1-\frac{1}{m}}}\bigg)^{\frac{2}{q'}-1}m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\\ &\le C(q)m_n\big(B_\rho\big)^{-\frac{2}{p}}\bigg(\frac{\theta_0}{2}\frac{\rho^2}{M^{1-\frac{1}{m}}}\bigg)^{\frac{2}{q'}-1}m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\\ &\le C(n,p,q)\bigg(\rho^{2\sigma_0}M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^j}{\omega}\Big)^2\theta_0^{\frac{2}{q'}}\bigg)\frac{1}{\theta_0\rho^2}\Big(\frac{\omega}{2^j}\Big)^2m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big),\end{aligned}$$

we obtain

$$I_2\le C(n,m,p,q)\bigg(\rho^{2\sigma_0}M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^j}{\omega}\Big)^2\theta_0^{\frac{2}{q'}}h(\rho,M,\omega)\bigg)\times\frac{1}{\theta_0\rho^2}\Big(\frac{\omega}{2^j}\Big)^2m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big).\tag{4.18}$$

Combining estimates (4.16), (4.17) and (4.18), we obtain

$$\begin{aligned}&\|\nabla(u-k)_+\|^2_{L^2(Q^{\theta_0}_{\rho_0,M})}\\ &\qquad\le C(n,m,p,q)\bigg(1+\rho^{2\sigma_0}M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^j}{\omega}\Big)^2\theta_0^{\frac{2}{q'}}h(\rho,M,\omega)\bigg)\frac{1}{\theta_0\rho^2}\Big(\frac{\omega}{2^j}\Big)^2m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\end{aligned}$$

and hence

$$\begin{aligned}&\Big(\frac{\omega}{2^{j+1}}\Big)^2m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\cap\{u>l\}\big)^2\\ &\qquad\le\frac{C(n,m,p,q)}{\theta_0^5}\Big(\frac{\omega}{2^j}\Big)^2\bigg(1+\rho^{2\sigma_0}M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^j}{\omega}\Big)^2\theta_0^{\frac{2}{q'}}h(\rho,M,\omega)\bigg)\\ &\qquad\qquad\times m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\cdot m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\cap\{k<u\le l\}\big).\end{aligned}$$

Summing over $i = r_0 + 1, \dots, q_0$, we have

$$\begin{aligned}
&\sum_{i=r_0+1}^{q_0} m_{n+1}\Big(Q^{\theta_0}_{\rho_0,M} \cap \Big\{u > \mu^+ - \frac{\omega}{2^{i+1}}\Big\}\Big)^2 \\
&\le \frac{C(n,m,p,q)}{\theta_0^5} m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big) \\
&\quad \times \sum_{i=r_0+1}^{q_0} \Big(1 + \rho^{2\sigma_0} M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^j}{\omega}\Big)^2 \theta_0^{\frac{2}{q'}} h(\rho, M, \omega)\Big) m_{n+1}\Big(Q^{\theta_0}_{\rho_0,M} \cap \Big\{\mu^+ - \frac{\omega}{2^i} < u \le \mu^+ - \frac{\omega}{2^{i+1}}\Big\}\Big) \\
&\le \frac{C(n,m,p,q)}{\theta_0^5} m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)\Big(1 + \rho^{2\sigma_0} M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^{q_0}}{\omega}\Big)^2 \theta_0^{\frac{2}{q'}} h(\rho, M, \omega)\Big) \\
&\quad \times \sum_{i=r_0+1}^{\infty} m_{n+1}\Big(Q^{\theta_0}_{\rho_0,M} \cap \Big\{\mu^+ - \frac{\omega}{2^i} < u \le \mu^+ - \frac{\omega}{2^{i+1}}\Big\}\Big) \\
&\le \frac{C(n,m,p,q)}{\theta_0^5} m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)^2\Big(1 + \rho^{2\sigma_0} M^{\frac{2}{q}(1-\frac{1}{m})}\Big(\frac{2^{q_0}}{\omega}\Big)^2 \theta_0^{\frac{2}{q'}} h(\rho, M, \omega)\Big).
\end{aligned}$$

We take $q_0 > 0$ enough large such that

$$\frac{2C(n,m,p,q)}{\theta_0^5(q_0 - r_0)} \le \nu^2.$$

Since

$$\sum_{i=r_0+1}^{q_0} m_{n+1}\Big(Q^{\theta_0}_{\rho_0,M} \cap \Big\{u > \mu^+ - \frac{\omega}{2^{i+1}}\Big\}\Big)^2 \ge (q_0 - r_0) m_{n+1}\Big(Q^{\theta_0}_{\rho_0,M} \cap \Big\{u > \mu^+ - \frac{\omega}{2^{q_0+1}}\Big\}\Big)^2,$$

we have

$$\begin{aligned}
m_{n+1}\Big(Q^{\theta_0}_{\rho_0,M} \cap \Big\{u > \mu^+ - \frac{\omega}{2^{q_0+1}}\Big\}\Big)^2 &\le \frac{2C(n,m,p,q)}{\theta_0^5(q_0 - r_0)} m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)^2 \\
&\le \nu^2 m_{n+1}\big(Q^{\theta_0}_{\rho_0,M}\big)^2
\end{aligned}$$

provided $\rho^{2\sigma_0} \le \min\{\theta_0^{-\frac{2}{q'}} 2^{-2q_0}, \delta_2^2\}\omega^2 M^{-\frac{2}{q}(1-\frac{1}{m})} h(\rho, M, \omega)^{-1}$, where $\delta_2 > 0$ is given by Lemma 4.3. Taking $\delta_1^2 := \min\{\theta_0^{-\frac{2}{q'}} 2^{-2q_0}, \delta_2^2\}$, we obtain Lemma 4.5. $\square$

Proof of Proposition 4.1. Let $0 < \nu < 1$ be chosen later. We take $\delta_1 > 0$ and q_0 as in Lemma 4.5. We introduce the following scale transform

$$\begin{aligned}
s = M^{1-\frac{1}{m}} t, \quad &\tilde{u}(s,x) = u(t,x), \quad \tilde{\eta}(s,x) = \eta(t,x), \\
\tilde{f}(s,x) = f(t,x), \quad &\tilde{g}(s,x) = g(t,x).
\end{aligned}$$

Then, using (2.2), we may rewrite the Caccioppoli estimate (4.9) as follows:

$$\begin{aligned}
&\sup_{s \in I^{\theta_0}_\rho} \int_{B_\rho} (\tilde{u}(s) - k)_+^2 \tilde{\eta}^2(s)\, dx + \iint_{Q^{\theta_0}_\rho} |\nabla(\tilde{u} - k)_+|^2 \tilde{\eta}^2\, dsdx \\
&\le C(m)\bigg\{\iint_{Q^{\theta_0}_\rho} (\tilde{u} - k)_+^2 \bigg\{\Big(\frac{M}{\mu^+}\Big)^{1-\frac{1}{m}} \partial_s \tilde{\eta}^2 + |\nabla \tilde{\eta}|^2\bigg\}\, dsdx \\
&\qquad + \tilde{h}(\rho, \omega)\Big(\int_{I^{\theta_0}_\rho} m_n\big(B_\rho \cap \{\tilde{u}(s) > k\}\big)^{q'(\frac{1}{2}-\frac{1}{p})}\, ds\Big)^{\frac{2}{q'}}\bigg\} \\
&\le C(m)\bigg\{\iint_{Q^{\theta_0}_\rho} (\tilde{u} - k)_+^2 \Big\{\partial_s \tilde{\eta}^2 + |\nabla \tilde{\eta}|^2\Big\}\, dsdx \\
&\qquad + \tilde{h}(\rho, \omega)\Big(\int_{I^{\theta_0}_\rho} m_n\big(B_\rho \cap \{\tilde{u}(s) > k\}\big)^{q'(\frac{1}{2}-\frac{1}{p})}\, ds\Big)^{\frac{2}{q'}}\bigg\}
\end{aligned} \tag{4.19}$$

where $\tilde{h}(\rho,\omega) := \|\tilde{f}\|^2_{L^q(L^p_{\mathrm{w}})(Q_\rho)} + \omega\|\tilde{g}\|_{L^{\frac{q}{2}}(L^{\frac{p}{2}}_{\mathrm{w}})(Q_\rho)}$.

We take $p_*, q_* > 0$ as in the proof of Proposition 3.1 and for $i \in \mathbb{N}$ we take $\rho = \rho_i,\ k = k_i,\ \tilde{\eta} = \tilde{\eta}_i$ satisfying $\tilde{\eta}_i \equiv 1$ on $Q^{\theta_0}_{\rho_{i+1}}$ and

$$\begin{aligned}
k_i &= \mu^+ - \frac{\omega}{2^{q_0}} + \frac{1}{2^{q_0+i+2}}\omega, \quad \rho_i = \frac{1}{2}\rho + \frac{1}{2^{i+2}}\rho,\\
Y_i &:= \frac{m_{n+1}\big(Q^{\theta_0}_{\rho_i} \cap \{\tilde{u} > k_i\}\big)}{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)},\\
Z_i &= \frac{\rho_0^2}{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}\bigg(\int_{I^{\theta_0}_{\rho_i}} m_n\Big(B_{\rho_i} \cap \{\tilde{u}(s) > k_i\}\Big)^{\frac{q_*}{p_*}}\,ds\bigg)^{\frac{2}{q_*}},\\
|\nabla\tilde{\eta}_i| &\le \frac{2}{\rho_i - \rho_{i+1}} \le \frac{12\cdot 2^i}{\rho_0}, \quad \partial_s\tilde{\eta}_i \le \frac{4}{\theta_0}\frac{1}{\rho_i^2 - \rho_{i+1}^2} \le \frac{48\cdot 2^{2i}}{\theta_0\rho^2}.
\end{aligned}$$

From (4.19) and $(\tilde{u} - k_i)_+ \le \frac{\omega}{2^{q_0+1}}$, we obtain

$$\begin{aligned}
&\|(\tilde{u} - k_i)_+\tilde{\eta}_i\|^2_{L^\infty(L^2)\cap L^2(\dot{H}^1)(Q^{\theta_0}_{\rho_i})} + \tilde{h}(\rho,\omega)\bigg(\int_{I^{\theta_0}_{\rho_i}} m_n\Big(B_\rho \cap \{\tilde{u}(s) > k_i\}\Big)^{q'(\frac{1}{2}-\frac{1}{p})}\,ds\bigg)^{\frac{2}{q'}}\\
&\le C(m)\bigg\{\Big(\frac{\omega}{2^{q_0}}\Big)^2\Big(\frac{1}{\theta_0} + 1\Big)\frac{2^{2i}}{\rho^2} m_{n+1}\Big(Q^{\theta_0}_{\rho_i} \cap \{\tilde{u} > k_i\}\Big)\\
&\quad + \tilde{h}(\rho,\omega)\bigg(\int_{I^{\theta_0}_{\rho_i}} m_n\Big(B_\rho \cap \{\tilde{u}(s) > k_i\}\Big)^{\frac{q_*}{p_*}}\,ds\bigg)^{\frac{2}{q_*}(1+\frac{2\sigma_0}{n})}\bigg\}\\
&\le C(m,\theta_0)\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}{\rho_0^2}\Big(\frac{\omega}{2^{q_0}}\Big)^2\\
&\quad \times\bigg\{2^{2i}Y_i + \tilde{h}(\rho,\omega)\Big(\frac{2^{q_0}}{\omega}\Big)^2\Big(\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}{\rho_0^2}\Big)^{\frac{2\sigma_0}{n}} Z_i^{1+\frac{2\sigma_0}{n}}\bigg\}.
\end{aligned}$$

Since $\delta_1 \le \theta_0^{-\frac{1}{q'}}2^{-q_0}$, we have

$$\tilde{h}(\rho,\omega)\Big(\frac{2^{q_0}}{\omega}\Big)^2\Big(\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}{\rho_0^2}\Big)^{\frac{2\sigma_0}{n}} \le C(n,p,q,\theta_0)$$

and hence

$$\|(\tilde{u} - k_i)_+\tilde{\eta}_i\|^2_{L^\infty(L^2)\cap L^2(\dot{H}^1)(Q^{\theta_0}_{\rho_i})} \le C(n,m,p,q,\theta_0)\Big(\frac{\omega}{2^{q_0}}\Big)^2\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}{\rho_0}\Big\{2^{2i}Y_i + Z_i^{1+\frac{2\sigma_0}{n}}\Big\}.$$

By the Ladyženskaja inequality (cf. Proposition A.1) and the Hölder inequality, we have

$$\begin{aligned}
\|(\tilde{u} - k_i)_+\tilde{\eta}_i\|^2_{L^2(Q^{\theta_0}_{\rho_i})} &\le \|(\tilde{u} - k_i)_+\tilde{\eta}_i\|^2_{L^{2+\frac{4}{n}}(Q^{\theta_0}_{\rho_i})}\|\chi_{\{\tilde{u}>k_i\}}\|^2_{L^{n+2}(Q^{\theta_0}_{\rho_i})}\\
&\le C(n,m,p,q,\theta_0)\Big(\frac{\omega}{2^{q_0}}\Big)^2\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)^{1+\frac{2}{n+2}}}{\rho_0^2}Y_i^{\frac{2}{n+2}}\Big\{2^{2i}Y_i + Z_i^{1+\frac{2\sigma_0}{n}}\Big\}
\end{aligned}$$

and

$$\|(\tilde{u} - k_i)_+\tilde{\eta}_i\|^2_{L^{q_*}(L^{p_*})(Q^{\theta_0}_{\rho_i})} \le C(n,m,p,q,\theta_0)\Big(\frac{\omega}{2^{q_0}}\Big)^2\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}{\rho_0^2}\Big\{2^{2i}Y_i + Z_i^{1+\frac{2\sigma_0}{n}}\Big\}.$$

Since

$$\begin{aligned}
\|(\tilde{u} - k_i)_+\tilde{\eta}_i\|^2_{L^2(Q^{\theta_0}_{\rho_i})} &\ge \|(\tilde{u} - k_i)_+\|^2_{L^2(Q^{\theta_0}_{\rho_{i+1}}\cap\{\tilde{u}>k_{i+1}\})}\\
&\ge (k_{i+1} - k_i)^2 m_{n+1}\Big(Q^{\theta_0}_{\rho_{i+1}} \cap \{\tilde{u} > k_{i+1}\}\Big)\\
&= \bigg(\frac{\omega}{2^{q_0+i+3}}\bigg)^2 m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)Y_{i+1}
\end{aligned}$$

and

$$\begin{aligned}\|(\tilde{u}-k_i)_+\tilde{\eta}_i\|^2_{L^{q*}(L^{p*})(Q^{\theta_0}_{\rho_i})} &\geq \|(\tilde{u}-k_i)_+\|^2_{L^{q*}L^{p*}(Q^{\theta_0}_{\rho_{i+1}}\cap\{\tilde{u}>k_{i+1}\})}\\ &\geq \left(\frac{\omega}{2^{q_0+i+3}}\right)^2\frac{m_{n+1}\big(Q^{\theta_0}_{\rho_0}\big)}{\rho_0^2}Z_{i+1},\end{aligned}$$

we obtain

$$Y_{i+1}\leq C(n,m,p,q,\theta_0)\Big\{2^{4i}Y_i^{1+\frac{2}{n+2}}+2^{2i}Y_i^{\frac{2}{n+2}}Z_i^{1+\frac{2\sigma_0}{n}}\Big\}$$

and

$$Z_{i+1}\leq C(n,m,p,q,\theta_0)\Big\{2^{4i}Y_i+2^{2i}Z_i^{1+\frac{2\sigma_0}{n}}\Big\}.$$

Considering the same calculation of (3.7) and (3.8), we obtain

$$Z_0\leq\begin{cases}C(n,p,q,\theta_0)Y_0^{\frac{2}{p*}} & \text{if}\quad q\geq p,\\[1ex] C(n,p,q,\theta_0)Y_0^{\frac{2}{q*}} & \text{if}\quad q<p.\end{cases}$$

Therefore, by Lemma A.4, there exists $0<\nu=\nu(n,m,p,q,\theta_0)<1$ such that if $Y_0\leq\nu$, then $Y_i\to 0$ as $i\to\infty$, i.e.

$$\tilde{u}(s,x)<\mu^+-\frac{\omega}{2^{q_0+2}}\quad\text{a.a. } (s,x)\in Q^{\theta_0}_{\frac{\rho}{2}}.$$

By Lemma 4.5, we obtain the upper bounds of u. □

APPENDIX A. APPENDIX

Their results are well-known, however we give the proof here for reader's convenience.

A.1. **Some Sobolev type inequality.**

Proposition A.1 (Ladyženskaja-Solonnikov-Ural'ceva [15, p.74])**.** *Let $I\subset\mathbb{R}$ be an open interval and let $\Omega\subset\mathbb{R}^n$ be a domain. Then for $f\in L^\infty(I;L^2(\Omega))\cap L^2(I;H^1_0(\Omega))$ and $p,q\geq 2$ satisfying*

$$\begin{aligned}&\frac{2}{q}+\frac{n}{p}=\frac{n}{2} & &\textit{if}\quad n\neq 2,\\ &\frac{2}{q}+\frac{n}{p}=\frac{n}{2}\quad\textit{without}\quad q=2,\ p=\infty & &\textit{if}\quad n=2,\end{aligned}$$

we obtain

$$\|f\|_{L^q(I;L^p(\Omega))}\leq C(n,p,q)(\|f\|_{L^\infty(I;L^2(\Omega))}+\|\nabla f\|_{L^2(I\times\Omega)}). \tag{A.1}$$

Proof. By the Gagliardo-Nirenberg-Sobolev inequality, we have

$$\|f(t)\|_{L^p(\Omega)}\leq C(n,p)\|\nabla f(t)\|^{\frac{n}{2}-\frac{n}{p}}_{L^2(\Omega)}\|f(t)\|^{1-(\frac{n}{2}-\frac{n}{p})}_{L^2(\Omega)}\quad\text{a.a. } t\in I.$$

Taking $L^q(I)$ norm on both side, we obtain (A.1). □

Proposition A.2 (Ladyženskaja-Solonnikov-Ural'ceva [15, p.91])**.** *Let f be a non-negative function belonging to $W^{1,1}(B_\rho)$ and let $l>k$. Then there exists a constant $C>0$ depending on n only such that*

$$(l-k)m_n\big(\{f>l\}\big)\leq\frac{C\rho^{n+1}}{m_n(B_\rho)-m_n\big(\{f>k\}\big)}\int_{\{k<f\leq l\}}|\nabla f|\,dx.$$

For the proof of Proposition A.2, we need the following Poincaré inequalities:

Lemma A.3 (Ladyženskaja-Solonnikov-Ural'ceva [15, Lemma 5.1 in p.89])**.** *Let $g\in W^{1,1}(B_\rho)$ ne a non-negative function and let $N_0:=\{g=0\}$. Let $\eta(x)=\eta(|x|)$ be a decreasing function of $|x|$ satisfying $0\leq\eta\leq 1$ and $\eta\big|_{N_0}\equiv 1$. Then for measurable set $N\subset B_\rho$, we have*

$$\int_N g(x)\eta(x)\,dx\leq\frac{C_n\rho^n}{m_n(N_0)}m_n(N)^{\frac{1}{n}}\int_{B_\rho}|\nabla g(x)|\eta(x)\,dx.$$

Proof. First we consider the case of $n \geq 2$. For $x \in N$, $x' \in N_0$, we have

$$g(x) = g(x) - g(x') = -\int_0^{|x'-x|} \frac{d}{dr} g(x + r\omega)\, dr \leq \int_0^{|x'-x|} |\nabla g(x + r\omega)|\, dr$$

where $\omega = \frac{x'-x}{|x'-x|}$. We now show

$$\eta(x) \leq \eta(x + r\omega) \quad \text{for} \quad 0 < r \leq |x' - x|. \tag{A.2}$$

Either if $|x| \leq |x'|$, then $x + r\omega \in B_{|x'|}$ by the convexity of $B_{|x'|}$. By the monotonicity of η, we have $\eta(x + r\omega) \geq \eta(x') = 1$. Otherwise, if $|x| > |x'|$, then $x + r\omega \in B_{|x|}$. Since $\eta(x + r\omega) \geq \eta(x)$, we obtain (A.2).

By (A.2), we have

$$g(x)\eta(x) \leq \int_0^{|x'-x|} |\nabla g(x + r\omega)|\eta(x + r\omega)\, dr.$$

Integrating over $x \in N$ and $x' \in N_0$, we have

$$m_n(N_0) \int_N g(x)\eta(x)\, dx \leq \int_N dx \int_{N_0} dx' \int_0^{|x'-x|} |\nabla g(x + r\omega)|\eta(x + r\omega)\, dr.$$

Let $g(x) = \eta(x) = 0$ on $x \in \mathbb{R}^n \setminus B_\rho$. Introducing the polar coordinate, we obtain

$$\begin{aligned}
&\int_{N_0} dx' \int_0^{|x'-x|} |\nabla g(x + r\omega)|\eta(x + r\omega)\, dr \\
&\leq \int_{B_{2\rho(x)}} dx' \int_0^{|x'-x|} |\nabla g(x + r\omega)|\eta(x + r\omega)\, dr \\
&\leq \int_{B_{2\rho(x)}} dx' \int_0^{|x'-x|} |\nabla g(x + r\omega)|\eta(x + r\omega)\, dr \\
&= \int_0^{2\rho} s^{n-1}\, ds \int_{\mathbb{S}^{n-1}} d\sigma \int_0^s \frac{|\nabla g(x + r\omega)|\eta(x + r\omega)}{r^{n-1}} r^{n-1}\, dr \quad (x' = s\sigma + x) \\
&= \int_0^{2\rho} s^{n-1}\, ds \int_{B_s(x)} \frac{|\nabla g(y)|\eta(y)}{|x - y|^{n-1}}\, dy \quad (y = x + r\sigma) \\
&\leq \frac{(2\rho)^n}{n} \int_{B_\rho} \frac{|\nabla g(y)|\eta(y)}{|x - y|^{n-1}}\, dy.
\end{aligned}$$

Therefore,

$$\begin{aligned}
m_n(N_0) \int_N g(x)\eta(x)\, dx &\leq \frac{(2\rho)^n}{n} \int_N dx \int_{B_\rho} \frac{|\nabla g(y)|\eta(y)}{|x - y|^{n-1}}\, dy \\
&= \frac{(2\rho)^n}{n} \int_{B_\rho} |\nabla g(y)|\eta(y)\, dy \int_N \frac{1}{|x - y|^{n-1}}\, dx.
\end{aligned}$$

We now show the following estimate:

$$\int_N \frac{1}{|x - y|^{n-1}}\, dx \leq (1 + \mathscr{H}^{n-1}(\mathbb{S}^{n-1})) m_n(N)^{\frac{1}{n}}, \tag{A.3}$$

where $\mathscr{H}^{n-1}(\mathbb{S}^{n-1})$ is the $(n-1)$-dimensional Hausdorff measure of the $(n-1)$-dimensional unit sphere. To show (A.3), let $\delta > 0$ to be chosen later. We split the integral

$$\begin{aligned}
&\int_N \frac{1}{|x - y|^{n-1}}\, dx \\
&\leq \int_{N \cap \{|x-y| \leq \delta\}} \frac{1}{|x - y|^{n-1}}\, dx + \int_{N \cap \{|x-y| \geq \delta\}} \frac{1}{|x - y|^{n-1}}\, dx =: I_1 + I_2.
\end{aligned}$$

By a simple calculation, we obtain

$$I_1 \le \int_0^\delta \frac{r^{n-1}}{r^{n-1}}\,dr \int_{\mathbb{S}^{n-1}} d\sigma = \delta \mathscr{H}^{n-1}(\mathbb{S}^{n-1}),$$
$$I_2 \le \int_N \frac{1}{\delta^{n-1}}\,dx \le \delta^{1-n} m_n(N).$$

Taking $\delta = m_n(N)^{\frac{1}{n}}$, we have $I_1 + I_2 \le (1 + \mathscr{H}^{n-1}(\mathbb{S}^{n-1}))m_n(N)^{\frac{1}{n}}$ and we obtain (A.3).

Using (A.3), we have

$$m_n(N_0) \int_N g(x)\eta(x)\,dx \le \frac{2^n(1 + \mathscr{H}^{n-1}(\mathbb{S}^{n-1}))}{n} \rho^n m_n(N)^{\frac{1}{n}} \int_{B_\rho} |\nabla g(y)|\eta(y)\,dy.$$

We consider the case $n = 1$. For $x \in N$ and $x' \in N_0$, we have

$$g(x) = g(x) - g(x') = \int_{x'}^x \frac{d}{dy} g(y)\,dy \le \left| \int_{x'}^x \left| \frac{d}{dy} g(y) \right| dy \right|.$$

Since

$$g(x)\eta(x) \le \left| \int_{x'}^x \left| \frac{d}{dy} g(y)\eta(y) \right| dy \right| \le \int_1^{-1} \left| \frac{d}{dy} g(y)\eta(y) \right| dy,$$

we obtain

$$\int_N g(x)\eta(x)\,dx \le m_n(N) \int_{-1}^1 |\nabla g(x)|\eta(x)\,dx.$$

□

Proof of Proposition A.2. Let

$$g(x) := \max\{l - k\,,\, (f - k)_+\} \in W^{1,1}(B_\rho), \quad N_0 := \{f < k\},$$
$$\eta(x) \equiv 1, \qquad N := \{f > l\}.$$

Then, by the Lemma A.3, we have

$$\int_N g(x)\,dx \le \frac{C_n \rho^n m_n\big(N\big)^{\frac{1}{n}}}{m_n\big(N_0\big)} \int_{B_\rho} |\nabla g(x)|\,dx,$$

hence

$$(l - k)m_n\big(\{f > l\}\big) \le \frac{C_n \rho^n m_n\big(\{f > l\}\big)^{\frac{1}{n}}}{m_n\big(\{f < k\}\big)} \int_{\{k < f \le l\}} |\nabla f(x)|\,dx.$$

□

A.2. **The recursive inequalities.**

Lemma A.4 (Ladyženskaja-Solonnikov-Ural'ceva [15, Lemma 5.7 in p.96])**.** *Let* $C, \varepsilon, \delta > 0$ *and* $b \ge 1$. *Assume that sequences* $\{Y_n\}_{n=0}^\infty$, $\{Z_n\}_{n=0}^\infty \subset (0, \infty)$ *satisfy*

$$Y_{n+1} \le Cb^n(Y_n^{1+\delta} + Y_n^\delta Z_n^{1+\varepsilon}),$$
$$Z_{n+1} \le Cb^n(Y_n + Z_n^{1+\varepsilon}). \tag{A.4}$$

Let

$$d := \min\left\{\delta, \frac{\varepsilon}{1+\varepsilon}\right\}, \ \lambda = \min\left\{(2C)^{-\frac{1}{\delta}} b^{-\frac{1}{\delta d}}\,,\, (2C)^{-\frac{1+\varepsilon}{\varepsilon}} b^{-\frac{1}{\varepsilon d}}\right\}.$$

Then, if $Y_0 \le \lambda$ *and* $Z_0 \le \lambda^{\frac{1}{1+\varepsilon}}$, *we obtain*

$$Y_n \le \lambda b^{-\frac{n}{d}}\,,\ Z_n \le (\lambda b^{-\frac{n}{d}})^{\frac{1}{1+\varepsilon}}. \tag{A.5}$$

Proof. Inequalities (A.5) are valid for $n = 0$. We prove (A.5) by induction. If (A.5) hold for n, then by (A.4), we have

$$Y_{n+1} \le 2C\lambda^{1+\delta} b^{n(1-\frac{1+\delta}{d})}\,,\ Z_{n+1} \le 2C\lambda b^{n(1-\frac{1}{d})}.$$

Since $\lambda \le (2C)^{-\frac{1}{\delta}} b^{-\frac{1}{\delta d}}$ and $d \le \delta$, we have

$$2C\lambda^{1+\delta} b^{n(1-\frac{1+\delta}{d})} \le \lambda b^{-\frac{1}{d}} b^{-\frac{n}{d}+n(1-\frac{\delta}{d})} \le \lambda b^{-\frac{n+1}{d}}.$$

Similarly, since $\lambda \le (2C)^{-\frac{1+\varepsilon}{\varepsilon}} b^{-\frac{1}{\varepsilon d}}$, we obtain

$$
\begin{aligned}
2C\lambda b^{n(1-\frac{1}{d})} &= 2C\lambda^{\frac{\varepsilon}{1+\varepsilon}} \lambda^{\frac{1}{1+\varepsilon}} b^{-\frac{n+1}{(1+\varepsilon)d}} b^{n(1-\frac{1}{d})+\frac{n+1}{(1+\varepsilon)d}} \\
&\le (\lambda b^{-\frac{n+1}{d}})^{\frac{1}{1+\varepsilon}} b^{n(1-\frac{\varepsilon}{(1+\varepsilon)d})}.
\end{aligned}
$$

Since $d \le \frac{\varepsilon}{1+\varepsilon}$, we find $1 - \frac{\varepsilon}{(1+\varepsilon)d} \le 0$ and hence we have (A.5) for $n+1$. □

A.3. **The weak L^p spaces and the Lorentz spaces.**

Let $\Omega \subset \mathbb{R}^n$ be a domain (not necessary bounded).

Definition A.5 (The Lorentz spaces)**.** For $1 \le p < \infty$, we define the Lorentz space $L^{p,\infty}(\Omega)$ by

$$L^{p,\infty}(\Omega) := \{ f \in L^1_{\mathrm{loc}}(\Omega) \; : \; \lambda^p \mu_{|f|}(\lambda) \text{ is bounded for all } \lambda > 0 \}$$

where $\mu_{|f|}(\lambda) := m_n\big(\{|f| > \lambda\}\big)$.

Proposition A.6 (cf. Benilan-Brezis-Crandall [4, pp.548])**.** *For* $1 < p < \infty$, *we have*

$$\frac{p-1}{p^{1+\frac{1}{p}}} \|f\|_{L^p_{\mathrm{w}}(\Omega)} \le \sup_{\lambda > 0} \lambda \mu_{|f|}(\lambda)^{\frac{1}{p}} \le \|f\|_{L^p_{\mathrm{w}}(\Omega)}.$$

Proof. First, we show $\sup_{\lambda>0} \lambda \mu_{|f|}(\lambda)^{\frac{1}{p}} \le \|f\|_{L^p_{\mathrm{w}}(\Omega)}$. For $\rho, \lambda > 0$, we take $K = \{|f| > \lambda\} \cap B_\rho$. Then we have

$$
\begin{aligned}
\|f\|_{L^p_{\mathrm{w}}(\Omega)} &\ge m_n\big(\{x \in \Omega \cap B_\rho : |f(x)| > \lambda\}\big)^{\frac{1}{p}-1} \int_{\{|f|>\lambda\}\cap B_\rho} |f(x)|\, dx \\
&\ge \lambda m_n\big(\{x \in \Omega \cap B_\rho : |f(x)| > \lambda\}\big)^{\frac{1}{p}}.
\end{aligned}
$$

Letting $\rho \to \infty$, we find

$$\lambda \mu_{|f|}(\lambda)^{\frac{1}{p}} \le \|f\|_{L^p_{\mathrm{w}}(\Omega)}.$$

Second, we show $\frac{p-1}{p^{1+\frac{1}{p}}} \|f\|_{L^p_{\mathrm{w}}(\Omega)} \le \sup_{\lambda>0} \lambda \mu_{|f|}(\lambda)^{\frac{1}{p}}$. We fix $\lambda_0 > 0$. For measurable set $K \subset \Omega$, we have

$$\int_K |f(x)|\, dx \le \lambda_0 m_n\big(K\big) + \int_{\{|f|>\lambda_0\}} |f(x)|\, dx.$$

By the above inequality, we have

$$
\begin{aligned}
\int_{\{|f|>\lambda_0\}} |f(x)|\, dx &= \int_0^\infty m_n\Big(\{x \in \{|f| > \lambda_0\} : |f(x)| > \lambda\}\Big)\, d\lambda \\
&= \int_0^{\lambda_0} m_n\big(\{|f| > \lambda_0\}\big)\, d\lambda + \int_{\lambda_0}^\infty m_n\big(\{|f| > \lambda\}\big)\, d\lambda \\
&= \lambda_0 \mu_{|f|}(\lambda_0) + \int_{\lambda_0}^\infty \mu_{|f|}(\lambda)\, d\lambda \\
&\le \lambda_0^{1-p} \sup_{\lambda>0} \lambda^p \mu_{|f|}(\lambda) + \sup_{\lambda>0} \lambda^p \mu_{|f|}(\lambda) \int_{\lambda_0}^\infty \lambda^{-p}\, d\lambda \\
&= \frac{p}{p-1} \lambda_0^{1-p} \sup_{\lambda>0} \lambda^p \mu_{|f|}(\lambda).
\end{aligned}
$$

Taking $\lambda_0^p m_n\big(K\big) = p \sup_{\lambda>0} \lambda^p \mu_{|f|}(\lambda)$, we find

$$\int_K |f(x)|\, dx \le \frac{p}{p-1} (p \sup_{\lambda>0} \lambda^p \mu_{|f|}(\lambda))^{\frac{1}{p}} |K|^{1-\frac{1}{p}}$$

or

$$\|f\|_{L^p_{\mathrm{w}}(\Omega)} \le \frac{p^{1+\frac{1}{p}}}{p-1} \sup_{\lambda>0} \lambda \mu_{|f|}(\lambda)^{\frac{1}{p}}.$$

□

Acknowledgments: The author would like to express his deepest gratitude to Professor Takayoshi Ogawa for his valuable comments and encouragement. The author also wishes to thank the referee for his valuable suggestions. This work is supported by the JSPS Research Fellowships for Young Scientists and the JSPS Grant-in-Aid for the JSPS fellows #21-1281.

REFERENCES

[1] Acerbi, E. and Mingione, G., *Gradient estimates for a class of parabolic systems*, Duke Math. J. **136** (2007), 285–320.

[2] Aronson, D. G. and Bénilan, P., *Régularité des solutions de l'équation des milieux poreux dans* $\mathbf{R}^N$, C. R. Acad. Sci. Paris Sér. A-B **288** (1979), A103–A105.

[3] Aronson, D. G. and Caffarelli, L. A., *The initial trace of a solution of the porous medium equation*, Trans. Amer. Math. Soc., **280** (1983), 351–366.

[4] Benilan, P., Brezis, H. and Crandall, M. G., *A semilinear equation in* $L^1(R^N)$, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **2** (1975), 523–555.

[5] Caffarelli, L. A. and Friedman, A., *Regularity of the free boundary of a gas flow in an* n*-dimensional porous medium*, Indiana Univ. Math. J. **29** (1980), 361–391.

[6] Caffarelli, L. A., Vázquez, J. L. and Wolanski, N. I., *Lipschitz continuity of solutions and interfaces of the* N*-dimensional porous medium equation*, Indiana Univ. Math. J., **36** (1987), 373–401.

[7] DiBenedetto, E., *Intrinsic Harnack type inequalities for solutions of certain degenerate parabolic equations*, Arch. Rational Mech. Anal. **100** (1988), 129–147.

[8] DiBenedetto, E., *Degenerate parabolic equations*, Universitext, Springer-Verlag, 1993.

[9] DiBenedetto, E., and Friedman, A., *Hölder estimates for nonlinear degenerate parabolic systems*, J. Reine Angew. Math. **357** (1985), 1–22.

[10] DiBenedetto, E., Gianazza, U. and Vespri, V., *Harnack estimates for quasi-linear degenerate parabolic differential equations*, Acta Math. **200** (2008), 181–209.

[11] DiBenedetto, E., Gianazza, U. and Vespri, V., *Harnack's inequality for degenerate and singular parabolic equations*, Springer Monographs in Mathematics, Springer, 2012.

[12] Duzaar, F. and Mingione, G., *Gradient estimates via non-linear potentials*, Amer. J. Math. **133** (2011), 1093–1149.

[13] Kinnunen, J. and Lewis, J. L., *Higher integrability for parabolic systems of* p*-Laplacian type*, Duke Math. J. **102** (2000), 253–271.

[14] Kuusi, T. and Mingione, G., *Nonlinear potential estimates in parabolic problems*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **22** (2011), 161–174.

[15] Ladyženskaja, O. A., Solonnikov, V. A. and Ural'ceva, N. N., *Linear and quasilinear equations of parabolic type*, Translations of Mathematical Monographs, Vol. 23, American Mathematical Society, 1967.

[16] Lions, J.-L., *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod, 1969.

[17] Luckhaus, S. and Sugiyama, Y., *Asymptotic propdf with the optimal convergence rate for a parabolic equation of chemotaxis in super-critical cases*, Indiana Univ. Math. J. **56** (2007), 1279–1297.

[18] Misawa, M., *Local Hölder regularity of gradients for evolutional* p*-Laplacian systems*, Ann. Mat. Pura Appl. (4) **181** (2002), 389–405.

[19] Oleĭnik, O. A., Kalašinkov, A. S. and Čžou, Y.-L., *The Cauchy problem and boundary problems for equations of the type of non-stationary filtration*, Izv. Akad. Nauk SSSR. Ser. Mat. **22** (1958), 667–704.

[20] Ogawa, T., *Asymptotic stability of a decaying solution to the Keller-Segel system of degenerate type*, Differential Integral Equations **21** (2008), 1113–1154.

[21] Ogawa, T. and Mizuno, M., *Regularity and asymptotic stability for the Keller-Segel system of degenerate type with critical non-linearity*, preprint.

[22] Ôtani, M., L^∞*-energy method and its applications*, Nonlinear partial differential equations and their applications, GAKUTO Internat. Ser. Math. Sci. Appl., vol. 20, Gakkōtosho, Tokyo, 2004, pp. 505–516.

[23] Porzio, M. M. and Vespri, V., *Hölder estimates for local solutions of some doubly nonlinear degenerate parabolic equations*, J. Differential Equations **103** (1993), 146–178.

[24] Sugiyama, Y. and Kunii, H., *Global existence and decay properties for a degenerate Keller-Segel model with a power factor in drift term*, J. Differential Equations **227** (2006), 333–364.

[25] Vázquez, J. L., *The porous medium equation*, Oxford Mathematical Monographs, The Clarendon Press Oxford University Press, 2007.

[26] Wiegner, M., *On* C_α*-regularity of the gradient of solutions of degenerate parabolic systems*, Ann. Mat. Pura Appl. (4) **145** (1986), 385–405.

(Masashi Mizuno) DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE AND TECHNOLOGY, NIHON UNIVERSITY, TOKYO 101-8308 JAPAN

E-mail address: `mizuno@math.cst.nihon-u.ac.jp`